

# Cool Subdwarf Investigations (CSI) I: New Thoughts for the Spectral Types of K and M Subdwarfs

Wei-Chun Jao<sup>1</sup>, Todd J. Henry<sup>1</sup>, Thomas D. Beaulieu<sup>1</sup>, John P. Subasavage<sup>1</sup>

*Department of Physics and Astronomy, Georgia State University, Atlanta, GA 30302-4106*

jao, thenry, beaulieu, subasavage@chara.gsu.edu

## ABSTRACT

Using new spectra of 88 K and M-type subdwarfs, we consider novel methods for assigning their spectral types and take steps toward developing a comprehensive spectral sequence for subdwarf types K3.0 to M6.0. The types are assigned based on the overall morphology of spectra covering 6000Å to 9000Å. The types and sequence presented link the spectral types of cool subdwarfs to their main sequence counterparts, with emphasis on the relatively opacity-free region from 8200–9000Å. When available, supporting abundance, kinematic, and trigonometric parallax information is used to provide more complete portraits of the observed subdwarfs. We find that the CaHn ( $n = 1-3$ ) and TiO5 indices often used for subdwarf spectral typing are affected in complicated ways by combinations of subdwarfs' temperatures, metallicities, and gravities, and we use model grids to evaluate the trends in all three parameters. Because of the complex interplay of these three characteristics, it is not possible to identify a star as an “extreme” subdwarf simply based on very low metallicity, and we suggest that the modifiers “extreme” or “ultra” only outline locations on spectroscopic indices plots, and should not be used to imply low or very low metallicity stars. In addition, we propose that “VI” be used to identify a star as a subdwarf, rather than the confusing “sd” prefix, which is also used for hot O and B subdwarfs that are unrelated to the cool subdwarfs discussed in this paper.

*Subject headings:* stars: subdwarfs — stars: abundances — stars: late-type — stars: classification — stars: fundamental parameters — stars: temperatures

---

<sup>1</sup>Visiting Astronomer, Cerro Tololo Inter-American Observatory. CTIO is operated by AURA, Inc. under contract to the National Science Foundation.

## 1. Introduction

The HR diagram is the most important map of stellar astronomy. It provides a relatively straightforward method for separating different stellar luminosity classes, e.g. supergiants, bright giants, giants, subgiants, main sequence dwarfs, and white dwarfs, using their colors and luminosities. However, spectroscopic and trigonometric parallax results have revealed a seventh distinct stellar luminosity class — the subdwarfs — that lie below the main sequence dwarfs on the HR diagram.

The realm of the subdwarfs has been previously explored by Sandage & Eggen (1959), Hartwick et al. (1984), Monet et al. (1992), Gizis (1997), and Jao et al. (2005), to name a few. Subdwarfs’ locations on the HR diagram are in part due to having metallicity abundances lower than most field stars, which causes their opacities to differ from those of regular dwarfs. Subdwarfs are sometimes called low metallicity halo stars or Galactic thick disk stars based on their spectroscopic features, kinematics, and/or ages (Digby et al. 2003; Lépine et al. 2003; Burgasser et al. 2003; Reid and Gizis 2005; Monteiro et al. 2006). Regardless of how they are described, subdwarfs are fundamentally different from their main sequence cousins.

Because of their generally high intrinsic velocities in the Galaxy, many subdwarfs have been selected using high proper motion efforts, such as the Lowell proper motion (Giclas et al. 1971, 1978), LHS (Luyten 1979), and Lépine-Shara Proper Motion-North (LSPM, Lépine & Shara 2005a) catalogs. Recently, subdwarfs have been selected by colors and spectroscopic observations via the Sloan Digital Sky Survey (SDSS, West et al. 2004). After initial flagging as a potential subdwarf, spectroscopic and astrometric (i.e. trigonometric parallax work) followup efforts are typically carried out to confirm or refute candidates as true subdwarfs. Past subdwarf identification efforts include Ryan and Norris (1991), Monet et al. (1992), Carney et al. (1994), Gizis (1997), Lépine et al. (2003), West et al. (2004), Reid and Gizis (2005) and Burgasser & Kirkpatrick (2006). Most of these studies confirmed subdwarfs spectroscopically, but only Monet et al. (1992) provides the crucial trigonometric parallaxes that allow subdwarfs to be placed on the HR diagram.

Gizis (1997) presented a pioneering effort to assign numeric subtypes for cool subdwarfs of spectral types K and M. First, he used the flux ratio of molecular band features, CaHn ( $n = 1-3$ ; we will use “CaH” to indicate all three bands or index values throughout this paper, unless otherwise specified) and TiO5 with pseudo-continuum points to calculate spectroscopic indices ( $f_{bands}/f_{continuum}$ ). On TiO5 vs CaH1 or TiO5 vs CaH2+CaH3 plots, two high order polynomial lines can be used to separate a continuous distribution of stars into three categories: regular dwarfs, subdwarfs, and extreme subdwarfs. Numerical subclasses were then assigned using two independent linear fits for subdwarfs and extreme subdwarfs.

For a decade, this methodology has been used to assign spectral types for cool subdwarfs.

However, the current method of assigning subdwarf spectral types is not directly linked to either their main sequence or giant counterparts, as is typically (but not always) the case with normal dwarfs and giants. In addition, we have found that many subdwarfs are assigned different subtypes even though their differences are limited to CaH. If these limitations can be overcome, a well-defined spectral sequence would benefit many research areas beyond classification efforts, including attempts to estimate effective temperatures and distances, as well as providing insight into understanding Galactic structure.

In this work, we first provide clarification and recommendations for subdwarf terminology by addressing the usage of the confusing spectral prefix “sd”. We then discuss our spectroscopic observations of 88 subdwarfs, generally targeting high proper motion stars in the southern sky. We next outline how synthetic spectra can assist us in understanding cool subdwarf spectral features. The bulk of this work describes a detailed effort to provide a subdwarf spectral sequence for stars having spectral types K3.0 through M6.0. We then apply this spectral typing method to those subdwarfs found in the SDSS database. Once we have a detailed understanding of what makes a star a subdwarf, we then discuss (1) why it is premature to assign precise parameters to subdwarf spectra, (2) misunderstandings related to the terms “extreme” and “ultra” subdwarfs, (3) why the old method works for dwarfs, but not subdwarfs, and finally (4) the recent set of subdwarf spectral standards from Lépine et al. (2007).

## 2. “VI” Subdwarfs are Different from “sd” Subdwarfs

The first subluminescent objects fainter than main sequence stars were reported by Adams & Joy (1922) when they were trying to determine the luminosities of A-type stars. In Adams et al. (1935) they called these stars “intermediate white dwarfs” to separate them from typical white dwarfs, and in the same paper reported the first six “intermediate white dwarfs” — now known as LHS 405 (sdF3), LHS 540 (F8IV), LHS 1501 (A4p), LHS 2194 (sdF5), HD 132475 (F5/F6V), and HIP 68321 (A4). All six have either A or F types (spectral types are from SIMBAD) in the modern MK spectral classification system. However, the term “subdwarfs” was not suggested until Kuiper (1939). He expected three classes of objects to be found in his spectroscopic survey of high proper motion stars— white dwarfs, stars of large spectroscopic parallaxes, and a class that was 2 or 3 magnitudes less luminous than main

sequence stars of the same color<sup>1</sup>. He suggested the name “subdwarfs” be used to represent this final, independent, class of stars. This name paralleled the use of the term “subgiants” to describe stars that fall below the giants on the HR diagram. The name also eliminated the confusion with white dwarfs, which are much less luminous than main sequence stars (and we now know are a completely different type of object). A year later, Kuiper (1940) reported the first three M-type subdwarfs — Kapteyn’s star, LHS 20, and LHS 64 (Gizis 1997 confirmed that all are M subdwarfs). Although they were termed subdwarfs, he used the same spectral classification as dwarfs (M0 and M2, see Kuiper 1940 Table 1). The “sd” spectral classification prefix for subdwarfs did not appear until Joy (1947), when he used the strengthening of the Lindblad depression around 4226Å.

In the late 1940’s through 60’s, the term “subdwarfs” also began being used for a class of underluminous blue stars (Humason & Zwicky 1947; Feige 1958; Greenstein 1966; Greenstein & Münch 1966). The terminology was based on the understanding that if these were high luminosity blue stars, their distances would be outside the Milky Way, so it was surmised that these stars should be underluminous, and therefore closer. Although their temperatures are similar to O and B dwarfs, their spectral features are, in fact, quite different. Generally, O-type subdwarfs (sdO) and B-type subdwarfs (sdB) both have broad Balmer absorption. sdBs have weak or no He lines, while sdOs have strong He II (4686Å) or other He II lines (Heber 1992). Since Feige (1958), such blue objects have been called sdO, sdB, or sdOB-type stars.

Thus, we are left with the unfortunate situation that there are two different classes of stars called “subdwarfs.” One is located at the cool end of the HR diagram while the other is at the hot end. The two classes of stars are subluminous for completely different astrophysical reasons but share the same “sd” spectral classification prefix. Cool subdwarfs usually have low metallicity (Chamberlain & Aller 1951; Greenstein & Eggen 1966; Mould 1976; Allard & Hauschildt 1995), so their opacities are different from dwarfs. For example, Allard & Hauschildt (1995) discussed the possible opacity sources for a solar type dwarf and a  $[m/H]=-2.5$  subdwarfs. TiO dominates the opacity sources in the optical band. However, because of a decreasing metallicity for subdwarfs, TiO opacity decreases dramatically. Hence, this less blanketing from TiO bands causes more continuum flux radiated from deeper and hotter layer of stellar atmosphere and their spectrum falls closer overall to that of a black-body, so these subdwarfs appear bluer than dwarfs, as shown in Figure 1<sup>2</sup>. To the contrary,

---

<sup>1</sup>Kuiper assumed all of these high proper motion stars had  $V_{tan}$  less than 474 km/sec. Under this assumption, he could then make crude estimates of their absolute magnitudes without having trigonometric parallaxes. The results placed some stars three magnitudes below the main sequence.

<sup>2</sup>The details of these synthetic spectra, the “GAIA model grids”, are discussed in section 4.

hot blue subdwarfs of types O and B are progenitors of white dwarfs, and their subluminous nature is not caused by metallicity at all. Instead, hot subdwarfs represent a stage in the stellar evolution cycle of an evolved star, and they happen to be crossing the main sequence at the moment of observation.

Because they *are* different kinds of stellar objects, we suggest that the two classes should not share the same spectral classification notation, “sd.” Roman (1955) argued that for types later than G0, the spectral notation “VI” should be used for stars that are  $\sim 1$ –2 magnitude less luminous than main sequence stars. Although Jaschek & Jaschek (1987) stated that “this designation (VI) should definitely be abandoned”, no specific reasons were actually given. Here we propose that the luminosity class “VI” should be adopted for cool subdwarfs, especially for K and M types. The three primary reasons are:

- Cool subdwarfs of types K and M lie clearly below main sequence stars on the HR diagram, forming an additional class of objects. Assigning their types as VI continues the progression outlined by other classes. Like the family of giants, which includes luminous supergiants (I), bright giants (II), normal giants (III), and subgiants (IV), the family of dwarfs includes main sequence dwarfs (V) and their subdwarf (VI) counterparts.
- Cool subdwarfs and OB subdwarfs are completely different types of objects with different origins. The notation “sd” is suitable for OB subdwarfs, which have not yet as a group been established to have any sort of sequence on the HR diagram, at least when using parallax measurements from the Yale Parallax Catalog, Hipparcos, and more recent efforts. On the other hand, there are dozens of cool subdwarfs with parallax measurements, and they do form a coherent group below the main sequence on the HR diagram, making a Roman numeral designation reasonable.
- Historically, the Roman numerals used for luminosity categorization track with different gravities for the stars classified, with higher gravities being assigned higher Roman numerals. We find that assigning “VI” for subdwarfs appropriately continues this trend. Figure 2 shows the mass-gravity relation using data from Table 1 of López-Morales (2007). In the best represented mass regime from  $0.35M_{\odot}$  to  $0.70M_{\odot}$ , a crude trend indicates that lower metallicity stars do indeed have higher gravities. Although much more data are needed to understand clearly how metallicity affects the mass-gravity relation, current evidence supports using the “VI” designation for low metallicity subdwarfs, which tend to exhibit higher gravities.

Thus, in order to separate the OB subdwarfs from the cool subdwarfs, we suggest the ‘sd’

prefix should not be used for low metallicity (and/or high gravity, as will be shown below) subdwarfs.

### 3. Observations and Reductions

#### 3.1. Observations

Our subdwarf targets were selected from several different efforts, including lists of spectroscopically identified subdwarfs (Gizis 1997; Reid and Gizis 2005), subdwarfs with parallax measurements (Jao et al. 2005; Costa et al. 2005, 2006), candidates from proper motion catalogs (Deacon et al. 2005; Subasavage et al. 2005a,b), and stars with metallicity measurements (Carney et al. 1994; Cayrel de Strobel et al. 2001; Nordström et al. 2004). We define our subdwarfs of interest to be those with  $V - K_s \geq 2.0$ ,  $[m/H] \leq -0.5$ , or having absolute magnitudes at least one magnitude less luminous in  $M_{K_s}$  than a fit to main sequence stars of comparable color with trigonometric parallaxes from the RECONS (Research Consortium on Nearby Stars) 10 pc sample (Henry et al. 2006).

Spectroscopic observations were made with the 1.5-m and 4.0-m telescopes at CTIO. For the observations on the 1.5-m from 2002 to 2006, the R-C spectrograph with a Loral 1200×800 CCD camera was used with the #32 grating (in first order) at tilt 15.1°. The order-blocking filter OG570 was utilized to provide spectra covering the range of 6000Å to 9500Å with a resolution of 8.6Å. The only variation in observing parameters was that during the May and December 2006 observing runs a larger slit width of 6" and 4", respectively, was used instead of the 2" slit used in previous runs, in order to minimize the differential color refraction (because the slit orientation was not changed during observations). For observations on the 4.0-m in 2002, the R-C spectrograph with a Loral 3K×1K CCD was used with the #181 grating (in first order) at tilt 58.8°. The order-blocking filter OG515 was utilized to provide spectra covering the range from 5500Å to 10000Å with a resolution of 6Å. Fringing at wavelengths longer than  $\sim 7000\text{\AA}$  in the 4.0-m data was removed by customized IDL routines. Bias frames and dome flats (and sky flats at the 1.5-m) were taken at the beginning of each night for calibration. At least two exposures were taken for each object to permit cosmic ray rejection. If stars were faint, additional observations were sometimes made. A 10 second Ne+He+Ar or Ne only arc lamp spectrum was recorded after each target to permit wavelength calibration. Several spectroscopic flux standard stars found in the *IRAF* spectroscopy reduction packages were observed during each observing run, usually nightly. Reductions were carried out in the standard way using IRAF reduction packages. Wavelength and flux calibrations were done using *onedspec.dispcor* and *onedspec.calibrate* within IRAF, respectively.

Many of the subdwarfs discussed in this paper have new trigonometric parallaxes and *VRI* photometry acquired during our southern nearby star program, CTIOPI (Cerro Tololo Inter-american Observatory Parallax Investigation, see Jao et al. 2005). In a future paper in this subdwarf series, we will present the astrometric (particularly trigonometric parallaxes) and photometric results.

### 3.2. Identifying Subdwarfs

During our five year spectroscopic campaign, we have acquired spectra for more than 900 objects. To glean subdwarfs from our spectroscopic database, we calculated their spectroscopic indices listed in Table 1 (targets listed alphabetically), mimicking the methodology outlined by Gizis (1997).

Figure 3 shows TiO5 plotted against CaH1 and CaH2+CaH3 for various samples of stars. Our subdwarfs are shown with solid circles. For comparison, small dots indicate main sequence stars from Hawley et al. (1996), while open triangles and squares represent subdwarfs and “extreme” subdwarfs from Gizis (1997) and Reid and Gizis (2005). Some stars (solid circles) having indices located near or in the main sequence regions in these plots have been manually checked to confirm that they are subdwarfs. Using the HR diagram in Figure 4, we confirm the low luminosities of our spectroscopically selected subdwarfs ( $V - K_s > 2.7$ ) that have accurate trigonometric parallaxes.

Although we focus primarily on the K and M subdwarfs for this study, we include a few G-type subdwarfs among those selected from metallicity measurements in the literature. For reference, we consider early K-type stars to have types K0 to K2, mid K-types to be K3 to K5, and late K-types to be K6 and later. We find that it is difficult, but possible, to separate late G from early K-type stars using our spectral coverage and resolution. Their continuum slopes have only slight differences across our wavelength window coverage, and there are no noticeable absorption differences beyond 7500Å. Spectra for G1V to K5V types from Jacoby et al. (1984) (resolution  $\sim 4\text{\AA}$ ) and Silva & Cornell (1992) (resolution  $\sim 11\text{\AA}$ ) are plotted in Figure 5. The only strong features are the absorption lines of Ba I (6497Å)<sup>3</sup> and H $\alpha$  (6563Å), with gradually increasing Ba I absorption and decreasing H $\alpha$  absorption as the effective temperature drops. These effects can be seen in both sequences, regardless of the spectral resolution. Our spectral resolution of 6–9Å falls between the resolution of

---

<sup>3</sup>Turnshek et al. (1985) noted that this feature at 6497Å is a blend of different atomic lines, including Fe I, Ba I, Ca I, Mn I, Co I, Ti I and II, and Ni I. Ba I is likely the dominant absorber because it has the largest Einstein coefficient in the NIST atomic spectra database.

the two sequences shown, so we can use the relative absorption strengths of Ba I and H $\alpha$  to separate G and K-type stars. In total, we have identified 88 K and M subdwarfs and five G type subdwarfs, using spectra with coverage from 6000Å to 9000Å.

### 3.3. Sorting Spectra

After reduction, the spectra were sorted into different bins based upon similarity in overall slope and features. This assured that stars in each bin had approximately the same temperature. However, several impediments to clean sorting were encountered:

- All mid K-type subdwarfs had spectra virtually indistinguishable from dwarf standard stars, yet they had low metallicity measurements and/or were found below the main sequence on the HR diagram (see top figure in Figure 6).
- Many subdwarf spectra placed into the same bin showed differences only in CaH (see middle figure in Figure 6).
- Many subdwarf spectra matched different dwarf standards at the blue ( $\lambda < 7570\text{\AA}$ ) and red ( $\lambda > 8200\text{\AA}$ ) ends (see bottom figure in Figure 6).

In order to understand what factors caused these anomalies, we next examine theoretical studies that provide synthetic spectra that can be compared directly to the observed spectra.

## 4. Grids of Synthetic Spectra

We use grids of synthetic spectra computed with PHOENIX codes (hereafter, GAIA model grids) to understand how subdwarfs’ physical parameters (temperature, metallicity and gravity) affect their spectra. The most recently released GAIA model grids (Brott & Hauschildt 2005) are available at an FTP site in Hamburg<sup>4</sup>. Gizis (1997), Woolf & Wallerstein (2005), and Burgasser & Kirkpatrick (2006) have all used these synthetic model grids to characterize subdwarfs, but an older version of the grids was used in all three cases. The version we employ here, 2.6.1, was released in late 2004. A comparison of one pre-2004 spectral model (provided by V. Woolf, private communication) and a new spectral model (from the GAIA model grids) for a cool subdwarf is shown in Figure 7. Improvements to the new models

---

<sup>4</sup><ftp://ftp.hs.uni-hamburg.de/pub/outgoing/phoenix/GAIA/v2.6.1/>



include (1) an enlarged and enhanced version of the equation of state, (2) more atomic, ionic, and molecular line opacities, (3) inclusion of the formation of dust particles for cool stars, and (4) microturbulence calculations. Additional water and TiO opacities and the inclusion of dust are enhancements particularly applicable to the low mass stars discussed here.

The two spectra shown in Figure 7 are virtually identical redward of 7000Å, but there are significant differences between 6500Å and 7000Å, where the CaH2 and CaH3 bands are found, and these differences will certainly affect evaluations done with the older models. The new model in this region has much shallower absorptions than the old one, which will affect metallicity and gravity estimates. A few narrow absorption features (Li I at 6103Å, Ca I at 6122Å and 6162Å) are also changed. As outlined in the discussion section §9, even the latest version of the model grids does not provide ideal matches to real spectra, so further progress can still be made.

#### 4.1. Identifying Mid K-type Subdwarfs

We use the latest GAIA model grids<sup>5</sup> to calculate predicted CaH band strengths and plot the derived indices against effective temperatures in Figure 8. We evaluate stars with  $T_{eff}$  between 2700K and 4500K and  $[m/H]$  between 0.0 and  $-3.0$ . For the moment, we adopt  $\log g = 5.0$  generically for subdwarfs. This gravity value does not apply to all types of subdwarfs, but we are presently interested in outlining the behavior of the CaH features with metallicity alone.

When  $T_{eff}$  is less than about 3300K, Figure 8 shows that the CaH1 index decreases (stronger absorption) when metallicity decreases from 0.0 to  $-2.0$  at fixed  $T_{eff}$ . However, the trend reverses for  $[m/H] = -2.5$  and  $-3.0$ . When  $T_{eff}$  is between 3300K and 3500K, there is a very weak relation between metallicity and the CaH1 index, but from 0.0 to  $-1.0$ , the relation (CaH1 index decreases when metallicity decreases) still holds. This relationship is degenerate for lower metallicities. For temperatures hotter than 3500K, the CaH1 index increases (weaker absorption) as metallicity decreases, in contrast to the low temperature region. For the CaH2+CaH3 index, the trends are generally the same, except that (1) at temperatures less than 3200K the index decreases (stronger absorption) only for 0.0 to  $-1.0$ , with a reversal for lower metallicities, and (2) the trend for higher temperature stars (increasing index, weaker absorption with lower metallicity) is the same, but the turnover is

---

<sup>5</sup>GAIA model grids also provide various values of  $\alpha$ -elements (O, Ne, Mg, Si, S, Ar, Ca, and Ti), which yield abundance ratios such as O/Fe, Ne/Fe, etc. Nissen & Schuster (1997) showed that most field halo stars have  $[\alpha/Fe]_* \approx [\alpha/Fe]_\odot$ , so we select  $[\alpha/\alpha_\odot] = 0.0$ .

near 3200K rather than 3500K.

Even more important than these subtleties is that overall, the hotter the subdwarf, the weaker its CaH bands. Note the collapse of any differences between indices for hotter stars of various metallicities in Figure 8. This collapse makes separating mid K-type subdwarfs from dwarfs based only on spectroscopic indices difficult using our spectral coverage (6000Å–9000Å) and resolution (6Å–8.6Å). An alternative method, such as the HR diagram shown in Figure 4, sufficiently solves the problem for mid K-type subdwarfs and will be discussed in the next section. Independent metallicity measurements via high resolution spectroscopic observations (Bonfils et al. 2005; Bean et al. 2006) can also be utilized. Our own spectra are not of sufficiently high resolution to measure metallicities, but all selected K-type subdwarfs have measured  $[m/H] \leq -0.5$  from other publications.

#### 4.2. Mid K-type Subdwarf Sample from Our Observations

There are 31 subdwarfs having CaH1 indices larger than 0.9 enclosed by the dashed box in Figure 3. In this region, there are no subdwarfs with previously measured CaH/TiO5 indices to compare to our new sample of mid K-type stars, making it difficult to separate the dwarfs and subdwarfs based on the CaH1 index alone. Fortunately, 21 of these 31 stars have trigonometric parallaxes, and are plotted with concentric circles in Figure 4. At least 15 of these stars are subdwarfs based on their locations one or more magnitudes below the main sequence on the HR diagram. The star with the largest offset is DEN0515-7211<sup>6</sup>, located at  $(V - K) = 3.3$ ,  $M_{Ks} = 9.7$ . This star is a full 4.5 magnitudes less luminous than the main sequence, but has no CaH or TiO5 features. Two additional stars (G016-009AB and G026-009ACD) above the main sequence are known to be double-line spectroscopic binaries with  $[m/H] < -0.5$ . These individual targets are discussed in section 6.2. The dwarf/subdwarf status of only four stars of the 21 remain ambiguous — we suspect that most of them are also subdwarfs, perhaps with as yet undetected companions elevating them into main sequence territory.

Generally, the spectroscopic index method fails to distinguish subdwarfs from dwarfs if the derived CaH1 index is greater than 0.9. With the benefit of additional trigonometric parallax information and/or metallicity measurements, however, we can conclude that nearly all of these mid K-type stars are indeed subdwarfs.

---

<sup>6</sup>This star was first reported in Costa et al. (2006) as reference star #4 in the LHS 1749 parallax field. We identify it henceforth as DEN0515-7211 (DENIS-P J051545.1–721122).

## 5. Late K-type and M-type Subdwarfs from GAIA Model Grids

For late K-type (redder than K5.0) and M-type stars, we use the GAIA model grids to understand how the effective temperatures, metallicities, and gravities affect the shapes and features of subdwarfs’ spectra. This analysis allows us to develop a spectral sequence for cool subdwarfs of types K6.0 to M6.0, which is tentatively extended blueward to K3.0 when additional information is incorporated. We plot GAIA synthetic noiseless spectra in Figure 9, at increments of 200K (cooler than 4000K) and 400K (hotter than 4000K) for stars with metallicities of  $[m/H] = 0.0, -1.0, \text{ and } -2.0$ . Because gravities have very limited impact on the overall shapes of the spectra (shown in the top panel of Figure 10), we do not show gravity plots with fixed temperatures and metallicities (of course, some features do change markedly with gravity, but not the overall slopes of the spectra). Based on the synthetic spectra from GAIA model grids, cool subdwarf spectra between 6000Å and 9000Å exhibit the following trends:

- The effects of metallicity are minimal in low resolution subdwarf spectra for stars with temperatures of 4400K and hotter. This makes it difficult to separate dwarfs and subdwarfs using low resolution spectra (as discussed in §4.1). However, from these noiseless spectra, we can still identify a few metallic lines showing metallicity trends. The most prominent feature is marked #1 at 6256Å in Figure 9. Unfortunately, we do not see this feature in any of our K-type spectra (nor is it listed in Table 1 of Turnshek et al. 1985), so we consider its validity questionable. The next prominent feature, marked #2 in Figure 9, is Ca I (6162Å). This line can possibly be used to distinguish subdwarfs from dwarfs (see examples discussed in §7.2), but in practice it is somewhat difficult to evaluate in real spectra (with noise) at our resolution.
- For stars with temperatures of 2800–4000K, metallicity strongly affects the spectra between 6000Å and 8200Å. This is the region that has been historically used to assign spectral types. In effect, subdwarfs with decreased metallicities have spectra that are “brightened” or “less blanketed” at the blue end, relative to solar metallicity stars. However, the continuum at wavelengths longer than 8200Å for temperatures 3400–4800K is nearly free of metallicity effects. We can therefore use the 8200–9000Å region to establish subtypes in the subdwarf spectral sequence because the slope is a function of temperature. This also allows us to mirror the spectral sequence for dwarfs, providing a useful link between the dwarf and subdwarf sequences.
- The TiO5 band strength at 7050–7150Å is very sensitive to metallicity for temperatures cooler than 4000K. As shown in the top panel of Figure 10, the TiO5 band strength is effectively independent of gravity. We can therefore use the TiO5 feature to separate

subdwarfs with different metallicities if their continua (8200–9000Å) are the same, regardless of their gravities.

- For a star of given temperature, stronger CaH bands could be caused by lower metallicity, as shown in Figure 9, or higher gravity, as shown in the top panel of Figure 10. If two subdwarfs have the same continua from 8200–9000Å and the same TiO5 band strength, but their CaH bands are different, we can rank them by their relative gravities.

Consequently, the impediments to sorting cool subdwarf spectra at our resolution discussed in §3.3 can be overcome by understanding the trends revealed in GAIA synthetic spectra: (1) the mid K-type dwarfs have the same spectra as dwarfs for our spectral coverage and resolution, (2) CaH features are affected by both metallicity and gravity, while the TiO5 band is affected by metallicity but not gravity, and (3) the continuum from 8200Å to 9000Å is not strongly affected by either metallicity or gravity, so can therefore be used for spectral sequencing.

## 6. Additional Evidence Supporting the Metallicity and Gravity Trends Indicated by the GAIA Models

To investigate the metallicity trends seen in GAIA models, we compare our available spectra to metallicity measurements provided independently by others. Measuring M dwarf metallicities is difficult, but several recent attempts have made progress (Valenti et al. 1998; Woolf & Wallerstein 2005; Bonfils et al. 2005; Bean et al. 2006). Six red dwarfs that we have observed are included in the study by Bonfils et al. (2005), comprising three pairs of M dwarfs of types M1.0V, M2.5V and M3.0V shown in Figure 11. Each pair includes a relatively low metallicity dwarf (gray), and a relatively high metallicity dwarf (black). The red ends (from 8200Å to 9000Å) of each pair match one another, but the blue ends of the lower metallicity members is brighter in each case, as predicted by the GAIA models. The effect is rather subtle, but the derived metallicities for each pair are not wildly different (none of the six stars are subdwarfs), and yet the trend is confirmed in all three cases. One caveat is that the metallicities from these six stars were determined from the polynomial relation in Bonfils et al. (2005), not measured directly from spectra, but this appears to be the best that can be done given the available data. We conclude that the metallicity trend revealed in the GAIA models (see Figure 9) is sound because it appears to be confirmed in real M dwarf spectra.

Contrary to the evidence for the metallicity trend, we have found no direct spectroscopic

results to support the gravity trend in dwarfs and subdwarfs. Direct gravity measurements are difficult because (1) stars must have both mass and radius measurements and (2) clean spectra without contamination from companions must be obtained. This limits the available target lists to eclipsing binaries such as those discussed in López-Morales (2007) with cleanly deconvolved spectra or visual binaries in which individual radii can be measured via long baseline interferometry. Neither class of objects yet provides a rich dataset for cool dwarfs or subdwarfs.

Hence, we rely on other observational or theoretical efforts to investigate the gravity trend. The top panel of Figure 10 indicates that other than some sharp metallic lines, CaH bands show the most prominent changes when gravity varies (TiO is unaffected by changing gravity). Öhman (1934) demonstrated that the CaH2 band is found in the spectra of M-type dwarfs, but is not observed in the spectra of M-type giants, thus identifying the CaH2 band as a gravity indicator to separate dwarfs and giants. In addition to the GAIA models, Mould (1976) also showed that for stars with effective temperatures of 3250K, a spectrum from his atmospheric model with  $\log g = 5.75$  has stronger CaH2 than a spectrum with  $\log g = 4.75$ . If the CaH2 band is a gravity indicator, we may presume that the same gravity effects for CaH1 and CaH3 bands will be seen.

As shown in the two panels of Figure 10, the GAIA models imply that the CaH band strengths are indicators of both gravity and metallicity differences. In reality, if two red subdwarfs have spectra with the same overall continua and slopes, as well as matching TiO band strengths (e.g. middle panel in Figure 6), the only remaining discrepancies will be at the CaH bands. We believe that such differences are caused by different gravities.

Obviously, there is not yet a wealth of accurate direct observational results of red dwarfs and subdwarfs that can be used to stress test the metallicity and gravity trends seen in GAIA models. Nonetheless, what little we do have supports the trends, so we use these trends to assist us in establishing the subdwarf spectral sequence discussed below.

## 7. Subdwarf Spectral Sequence

### 7.1. Procedures

In addition to presenting 88 subdwarfs, a goal of this project is to establish a subdwarf spectral sequence that mirrors the sequence for dwarfs and considers the latest available synthetic models. Although synthetic models are not yet capable of fully representing the complicated spectra of these cool stars, the models can be used to investigate the primary factors that affect subdwarfs’ spectra. Specifically, we examine the spectra in the frame-

work of what appear to be the three main drivers of the trends observed — temperature, metallicity, and gravity.

We first separate our available subdwarf spectra into several groups that have similar overall slopes. Known subdwarfs such as LHS 12, GJ 161, and LHS 2734A are used as anchor points. Within each group, several stars that have high surface gravities are obvious because their spectra match except in CaH, implying nearly identical temperatures and metallicities.

We then compare each subdwarf’s spectrum with our sequence of dwarf spectral standards<sup>7</sup> that span types from K0.0 to M9.0 using an IDL program to find the closest match to the continuum slope in the region 8200–9000Å. Visual checks of the matches between all subdwarf and standard dwarf spectra are also made to ensure match quality.

## 7.2. Results and Notes on Objects

Based on the 88 confirmed cool subdwarf spectra we have, we present a sequence of subdwarfs with spectral types spanning K3.0 to M6.0, listed in Table 2. We also identify five additional G type subdwarfs. After the five G-type subdwarfs, we sort the cool subdwarfs from K3.0[VI] to M6.0VI, using double lines in Table 2 to separate each type.

As discussed in §4.1, because mid K subdwarfs are virtually indistinguishable from K dwarfs at our spectral resolution, we use [VI] to indicate their questionable luminosity classes, which are currently based on their metallicities, kinematics, or locations on the HR diagram. We anticipate that higher resolution spectra will reveal these stars to be subdwarfs. A colon after the type indicates that we have had difficulty in assigning a subtype, metallicity, or gravity.

---

<sup>7</sup>Our dwarf standard sequence is a hybrid of cool dwarf standards from Gray et al. (2006) for K stars, and Boeshaar (1976) and Kirkpatrick et al. (1991) for M stars. The dwarf standard sequence is the topic of a future publication (Beaulieu et al. 2009). Our spectral sequencing efforts began before the recent spectral sequence from SDSS (Bochanski et al. 2007) was released. We continue to use our standard sequence because (1) stars in our sequence have trigonometric parallaxes so we can understand how metallicities and gravities affect stars’ positions on the HR diagram, (2) the SDSS sequence does not include K-type stars, (3) we have acquired spectra for many K-type stars from Gray et al. (2003) and Gray et al. (2006) so we have benchmark K dwarf spectral standards to link to the M dwarf sequence, (4) SDSS standard stars have telluric lines removed (It is difficult for us to make comparisons to SDSS spectra in the 6800–7100Å region, which includes O<sub>2</sub> and H<sub>2</sub>O absorption. This region overlaps with the CaH3 and TiO5 bands. Thus, all subdwarfs would appear to have high gravity when compared to the SDSS standards.), and (5) our dwarf spectra have been acquired using the same telescope/instrument/observing protocols as used for the subdwarfs, and the data have been reduced identically, so they are systematically consistent.

Within each type for which sufficient spectra are available, we sort targets by their metallicities (lowest metallicity first). A letter “m” indicates a star having the same metallicity as a main sequence dwarf, while more negative signs indicate lower metallicities, e.g. m— is more metal poor than m–. We use as many as six negative signs, because in the case of the M1.0VI type, we have 23 stars that fall into seven different metallicity categories. Gravities are indicated by “g” with additional plus signs for higher gravities, e.g. g++ is higher gravity than g+. A baseline subdwarf of low metallicity is assigned m– and g. If a subdwarf has a similar metallicity but higher gravity, the designations are m– and g+. A few stars appear to have solar metallicity and are subluminescent only because of high gravities; these stars are assigned m and g+. We have removed any stars from this study that might have gravities lower than main sequence stars, i.e. slightly evolved stars such as subgiants, that would have gravity g–.

Note that values for metallicity and gravity are not comparable across all spectral types, i.e. M1.0VI with metallicity “m–” is not equivalent to M2.0VI with metallicity “m–,” nor is “g+” for M1.0VI the same as “g+” for M2.0VI. We can hope to formalize a definitive subdwarf spectral sequence that includes temperature, metallicity, and gravity trends when hundreds of systematically consistent subdwarf spectra and improved model grids are available. Objects with virtually identical spectra are listed in alphabetical and/or numerical order.

We discuss each spectral subtype from K3.0[VI] to M6.0VI in the following sections and highlight noteworthy subdwarfs of various types. The order of the highlighted targets is based on their metallicities or gravities. Colored spectra in the Figures in these sections are illustrative only — they do not represent any numeric metallicities or gravities, so a red-lined spectrum in one type does not necessarily have similar attributes to a red-lined spectrum in another type.

### 7.2.1. K3.0[VI] to K5.0[VI] types

We begin with K3.0[VI], for which it is still difficult to separate subdwarfs from dwarfs at our spectral resolution. Figure 9 shows how similar mid K-type subdwarfs and dwarfs are (temperatures 4400K and 4800K). For these stars, we use other independent published measurements, e.g. [m/H] values, to confirm their subdwarf natures. We currently call stars with  $[m/H] \leq -0.5$  subdwarfs. The K3.0[VI] to K5.0[VI] spectra are shown in Figure 12.

**G 016–009AB (K3.0[VI])** Goldberg et al. (2002) reported this star to be a double-lined spectroscopic binary with  $P=9.9$  days,  $T_{eff} = 5903K$ , and having a mass ratio ( $M_1/M_2$ )

of  $\sim 1.22$ . Given that the components are presumably coeval, the combined spectrum should represent the metallicity for each component. Although their estimated temperature is that of a G type star, our spectrum has the same slope as a K3.0V. Cayrel de Strobel et al. (2001) and Goldberg et al. (2002) reported measurements of  $[\text{Fe}/\text{H}] = -0.7$  and  $-1.0$  respectively, while Laird et al. (1988) found  $[\text{m}/\text{H}] = -1.11$ . Apparently, it is a low metallicity star and its location on the HR diagram is elevated because of multiplicity. This system is also reported to be a photometrically variable system in Kazarovets et al. (2006). This system demonstrates that low metallicity subdwarfs do not have different spectra from their main sequence counterparts at our wavelength coverage and resolution, so we assign its luminosity class as [VI] as a result its low metallicity measurements.

**LHS 2467 (K4.0[VI])** Our spectrum shows that it is a K4.0[VI], rather than a G7V type, as reported in Bidelman (1985). Its continuum is not as blue as K0.0V and its Ba I/H $\alpha$  lines are not as weak/strong as shown in Figure 5 for a G type star. Its weighted mean parallax from ESA (1997) and van Altena et al. (1995) is  $10.63 \pm 1.88$  mas and its proper motion is  $0''.98 \text{ yr}^{-1}$  (Luyten 1979), indicating  $V_{\text{tan}} = 437 \text{ km sec}^{-1}$ . Ryan and Norris (1991) reported the star to have  $V_{\text{rad}} = 44 \text{ km sec}^{-1}$ . After removing the solar motion, LHS 2467 has  $(U, V, W) = (281.4, -304.8, -41.9) \text{ km sec}^{-1}$ . The extremely high tangential and space velocities are indicative of a subdwarf.

**G 026–009ACD (K5.0[VI])** The wide ( $132''$ , corresponding to 6520 AU at the system’s distance of 49.4 pc) common proper motion companion, B, in this quadruple system is a white dwarf known as G026–010 or G026–009B, that during its planetary nebula phase should have had limited impact on the metallicity of the close ACD triple. Peterson et al. (1980) reported G026-009AC to be double-lined spectroscopic binary with  $P=3.75$  days, with a mass ratio  $(M_1/M_2) \sim 1.25$ . Allen et al. (2000) reported a third component, D,  $0''.7$  away from G026-009AC. Morrison et al. (2003) and Allen et al. (2000) both report the system (ACD) to have low metallicity, measuring  $[\text{Fe}/\text{H}] = -0.91$  and  $-1.19$ , respectively. Both the  $V$  and  $K_s$  magnitudes include three stars, causing the point to be improperly placed on the HR diagram.

**G 022-015 (K5.0[VI])** Woolf & Wallerstein (2005) reported this system to have  $[\text{Fe}/\text{H}] = -0.61$ . G022-015’s spectrum is virtually identical to G026-009ACD, as shown by plotting both together in Figure 12.

**GJ 223.1 (K5.0[VI])** This star has the same slope as the K5.0V standard and its location on the HR diagram is on the main sequence line. Nonetheless, Woolf & Wallerstein (2005) reported it to have  $[\text{m}/\text{H}] = -0.62$ , so we tentatively consider it to be a subdwarf until further information indicates otherwise.



Although not definitive, the Ca I line at  $6162\text{\AA}$  in these spectra seems to show the metallicity trend seen in the noiseless model spectra plotted in Figure 9 and discussed in section 5. LHS 2467 has a much weaker Ca I line, and G016–009AB and G026–009AB have slightly weaker Ca I lines than seen in the dwarfs’ spectra. However, G022–015 and GJ223.1 both have approximately the same Ca I line strengths as dwarfs. Contrary to the Ca I line, the Ca II lines at ( $8542\text{\AA}$  and  $8662\text{\AA}$ ) do not show any metallicity trend from our spectra. Because these prominent lines fail to separate subdwarfs from dwarfs at our resolution and S/N, other evidence, i.e. kinematics, parallaxes or independent metallicity measurements, are required to identify the subdwarfs. An alternative method to separate the subdwarfs from the dwarfs is to obtain spectra that include the MgH bands at  $4845\text{\AA}$ ,  $5211\text{\AA}$ , and  $5621\text{\AA}$  that Bessell (1982) pointed are sensitive to metal abundance for temperatures hotter than  $4500\text{K}$ .

Several possibilities may explain the locations of the three subdwarfs, LHS 2467, G 022-015, and GJ 223.1 on the HR diagram just below or on the main sequence line: (1) they are subdwarfs with unseen companions that brighten their  $M_{K_s}$  magnitudes, (2) they are slightly evolved and have moved significantly from the subdwarf zero age main sequence line, or (3) their metallicity measurements are not accurate and they are, in fact, main sequence dwarfs.

**LHS 125, LHS 232 and LHS 327** The spectra for these three stars, shown in Figure 13, match neither the LHS 2467 K4.0[VI] spectrum nor a K4.0V spectrum, although the spectra have the overall slopes and the stars have  $V - K_s$  colors indicative of K4.0 stars. They are tentatively assigned types of K4.0[VI] based on their locations on the HR diagram.

### 7.2.2. *K6.0VI*

We identify four stars to have spectral type K6.0VI, shown in Figure 14. Their spectral slopes fall between K5.0V and K7.0V.

**LHS 193A** This is a binary with a separation of  $12''.59$  comprised of a cool subdwarf (A) and a DC-type white dwarf (B) with a featureless spectrum (Monteiro et al. 2006). At the system’s distance of 31.2 pc, the large separation of the pair corresponds to 393 AU, indicating that significant pollution of the subdwarf by the white dwarf during its planetary nebula phase seems unlikely. From Figure 14, it is clear that LHS 193A’s spectral slope is between K5.0V and K7.0V. Although the spectrum is a near match to K5.0V between  $8200\text{\AA}$  and  $9000\text{\AA}$ , the blue end ( $6000\text{\AA}$ – $7500\text{\AA}$ ) is too low to be a K5.0V (as a subdwarf, the blue end of the spectrum would have to be above the dwarf standard). As shown in

Figure 9, for a star with 4400K (a K5.0V from Allen’s Astrophysical Quantities 2000), there is effectively no difference in the continua of spectra for dwarfs and subdwarfs. Hence, we assign LHS 193A a type of K6.0VI because its TiO5 band is “brightened” (low metallicity), as shown in Figure 9. Given its low metallicity, location on the HR diagram (shown in the inset of Figure 14), its  $V_{tan} = 147.3 \text{ km sec}^{-1}$  (Jao et al. 2005), and its age of 6–9 Gyr based on its white dwarf companion’s cooling age (Monteiro et al. 2006), it is most likely a thick disk subdwarf.

**LHS 73 and LHS 227** These two objects have virtually identical spectra, so we only show LHS 227 in Figure 14 for clarity. The only differences between these spectra and that of LHS 193A are in CaH. Using results from the top panel of Figure 10, we find that LHS 73 and LHS 227 have higher gravities than LHS 193A.

We have acquired spectra for both LHS 72 and LHS 73, a wide ( $\sim 96''$  separation) common proper motion pair. Unfortunately, the spectrum of LHS 72 ( $V - K_s = 3.29$ ,  $M_{K_s} = 6.69$ ) is poor and requires re-observation, although it would fall in the subdwarf region of the HR diagram. LHS 73 has  $V - K_s = 3.42$  and  $M_{K_s} = 7.27$  and is plotted in Figure 14, also clearly in the subdwarf region. Rodgers & Eggen (1974) previously identified both objects as subdwarfs, but no spectral types were given. Bidelman (1985) identified LHS 72 and LHS 73 as K4 and K5 dwarfs, respectively. Reyl   et al. (2006) reported LHS 72/73 (they identified LHS 73 as sdK7) to be the nearest subdwarf binary based on their spectroscopic parallax (18.7 pc), although the trigonometric parallax from YPC is  $37.6 \pm 8.9 \text{ mas}$  ( $21.5 \text{ pc} < d < 34.8 \text{ pc}$ ). The nearest known subdwarf binary system is actually  $\mu \text{ Cas AB}$ , at a distance of 7.5 pc ( $\pi_{trig} = 132.4 \pm 0.6 \text{ mas}$ , ESA 1997).

**LHS 161** This star was previously reported to be an esdM2.0 extreme subdwarf (Gizis 1997). However, its slope and spectral features, like LHS 227, are too hot for type M2.0VI. As shown in Figure 14, LHS 161 has stronger CaH than LHS 193A and LHS 227, but the rest of the spectrum is the same. We conclude that LHS 161 has higher gravity than the other two subdwarfs. LHS 161’s very strong CaH lines and consequent CaH and TiO5 indices place it in the previously called “extreme” subdwarf region, which typically implies that the star has very low metallicity. It appears, from our current understanding, that the strong CaH in LHS 161 may not be linked to metallicity, because the TiO5 band is not significantly different from LHS 193A, LHS 73 or LHS 227.

In the inset of Figure 14 we connect LHS 193A, LHS 73, LHS 227, and LHS 161 on the HR diagram with thick arrows to outline a sequence of increasing gravity effects that shift their locations to the lower right (less luminous and redder). We will soon see more such gravity effects, which are always towards lower luminosities.

### 7.2.3. K7.0VI

We identify four stars to have spectral type K7.0VI, shown in Figure 15. This is the earliest spectral type for which we see clear metallicity effects in the blue regions of our spectra.

**LHS 2734A** This is the primary in a new common proper motion system discovered during CTIOPI, for which we measure the components to have  $\mu = 0''.59$  and  $0''.60 \text{ yr}^{-1}$  at position angles  $228.2^\circ$  and  $228.4^\circ$  for the A and B components, respectively. The B component is located  $68''.8$  away at position angle  $162.4^\circ$ . Unfortunately, both components are beyond our current trigonometric parallax limit (100 pc), but the A component provides a useful subdwarf anchor point even though its location on the HR diagram is unknown.

There are three reasons to adopt LHS 2734A as a reliable subdwarf anchor. First, we measure a zero parallax, implying that its  $V_{tan}$  is at least 280 km/sec (assuming a distance of 100 pc). A tangential velocity of this size is indicative of a subdwarf. Second, LHS 2734B ( $V - K_s = 3.90$ ), is redder than LHS 2734A ( $V - K_s = 3.23$ ), and later type subdwarfs are more easily separated from dwarfs, as shown in Figure 9. As discussed in §7.2.6, LHS 2734B is clearly a subdwarf. Assuming this common proper motion pair formed at the same time with similar “genetics”, both components are subdwarfs. Third, Figure 15 shows the comparison between K7.0V and LHS 2734A, in which the blue end of LHS 2734A’s spectrum is clearly “brightened”. This indicates that LHS 2734A has lower metallicity than K7.0V, as expected for a subdwarf.

**LHS 164** This star is four full magnitudes below the main sequence line in the inset in Figure 15. We find a good match at the red end between our K7.0V standard and LHS 164’s spectrum, while the blue end is elevated. We conclude that LHS 164’s metallicity is even lower than LHS 2734A’s.

Thus, our K7.0V standard, LHS 2734A, and LHS 164 form a sequence of decreasing metallicity. The CaH and TiO5 band depths significantly decrease as the metallicity drops, matching the effects discussed in section 4.1 and Figure 8.

**DEN0515–7211 and SCR 0708–4709** These two spectra are virtually identical, and are assigned type K7.0VI: (Figure 15) because the spectra do not match either the K7.0V standard, LHS 164, or LHS 2734A. DEN0515–7211 is nearly five magnitudes less luminous in  $M_{K_s}$  than the main sequence line, so it is certainly a subdwarf. SCR 0708–4709 ( $\mu = 0''.402 \text{ yr}^{-1}$ , Subasavage et al. 2005a) has almost the same spectrum as DEN0515–7211 and falls in the subdwarf region on the reduced proper motion diagram (see Subasavage et al. 2005a).

#### 7.2.4. *M0.0VI*

We show four stars with spectral type M0.0VI in Figure 16. This is the first group in this manuscript for which we see clear effects of both metallicity and gravity.

**LHS 244, LHS 418, and LHS 165** These three spectra have the same continua at the red end but are very different at the blue end. The differences are caused by metallicity, as seen in Figure 9, with the main sequence standard–LHS 244–LHS 418–LHS 165 trending to lower metallicities. The blue ends of the subdwarf spectra have clearly “brightened” along this sequence, which is consistent with the theoretical models. In addition, as discussed in §4.1, CaH1 absorption decreases (increasing index) as metallicity decreases if  $T_{eff}$  is greater than 3500K. LHS 165 is redder in  $V - K_s$  and has stronger CaH1 absorption than LHS 418, so we suspect LHS 165 has a higher gravity (similar to gravity effects shown in Figure 14). We also find that LHS 418 has weaker CaH1 absorption than LHS 244, indicating that LHS 244 could also possibly have higher gravity than LHS 418. LHS 244 is much redder than LHS 418 and LHS 165 in the HR diagram is because of a combination of gravity and metallicity effects.

**LHS 424** This star has the same spectrum and color as LHS 418, so we assign the same spectral type. We note that LHS 424 is a full magnitude less luminous in  $M_{Ks}$ , perhaps hinting that LHS 418 is a multiple system.

**LHS 300AB** This is a close binary with a separation of  $\sim 4''.3$  (Jao et al. 2003) comprised of a cool subdwarf (A) and a DC-type white dwarf (B) with a featureless spectrum (Monteiro et al. 2006). The spectrum obtained includes both components, but with  $\Delta VRI = 4.61, 4.85, 4.96$  mag (Monteiro et al. 2006), the contamination of the cool subdwarf spectrum from the white dwarf is negligible. At a distance of 31.0 pc, the projected separation implies a distance of 133 AU between the two components, indicating that the subdwarf’s composition was unlikely to be significantly contaminated by the evolved star during its planetary nebula phase. The spectrum does not match other M0.0VI stars at the blue end, but it matches the red end of M0.0V and M0.0VI well. We therefore assign it a type of M0.0VI:.

#### 7.2.5. *M0.5VI*

This is the first half spectral type we currently assign for subdwarfs. We see both metallicity and gravity effects for M0.5VI, as shown in Figures 17 and 18.

**LHS 507, SCR 0709–4648, and LHS 12** These three stars, along with our M0.5V standard, illustrate a nice four-step trend in metallicity. With the lowest metallicity, LHS 507 is significantly bluer than LHS 12, which is, in turn, bluer than our main sequence standard. We do not yet have a trigonometric parallax for SCR 0709–4648, so we cannot confirm its location on the HR diagram. LHS 12 was previously reported to be a type sdM0.0 in Gizis (1997).

**LHS 401** This star has stronger CaH bands and therefore higher gravity than LHS 507, which has extremely weak CaH1. LHS 401 is redder than LHS 507, as expected, but is slightly more luminous, which is not consistent with the gravity effects seen for types K6.0VI and M0.0VI. We believe this inconsistency is a result of the large absolute magnitude errors for these two objects.

**LHS 521** This star has a spectrum nearly identical to LHS 507, except at 6000–6200Å. Its CaH2 feature is slightly different from LHS 507, so we assign it a type of M0.5VI:.

**LHS 367 and LHS 299** Both objects have trigonometric parallaxes. Their spectra are similar to SCR 0709–4648 and LHS 12, respectively (shown in Figure 18), so they also have similar offsets in metallicities from one another. We assign them as M0.5VI: because of (1) discrepancies at 6000Å–6200Å, (2) LHS 299 has deeper CaH1 absorption than LHS 12 and different CaH2 absorption, and (3) their locations on the HR diagram are redder than their respective comparison stars. LHS 299 may have a higher gravity than LHS 12, but a smaller absolute magnitude error is needed to confirm this speculation.

**LHS 360** This star appears to have higher gravity than LHS 12 because the CaH features are deeper, but its location on the HR diagram does not match its stronger gravity because its  $V - K_s$  color is slightly bluer than LHS 12. Costa et al. (2006) reported that LHS 360 has  $V_{tan} \sim 524 \text{ km s}^{-1}$ . Based on its very high  $V_{tan}$  and location on the HR diagram, we assign it a type of M0.5VI:.

#### 7.2.6. *M1.0VI*

The M1.0VI type has the largest number (23) of confirmed subdwarfs in our current sample. The spectra of 16 different M1.0VI stars are shown in Figures 19 and 20. Many of these stars have similar colors, as identified in a reduced proper motion diagram derived using results from our SuperCOSMOS-RECONS (SCR) survey (Subasavage et al. 2005a,b). Given the rich dataset for this type, we first discuss the sample in terms of metallicity and gravity effects, then discuss individual targets.

*Metallicity effects* Perhaps better than for any other spectral type available, the M1.0VI stars show a beautiful trend in metallicity in their spectra, as shown in the top of Figure 19. The red ends of the spectra match the M1.0V standard spectrum, but the blue ends of the spectra are very different because of metallicity effects. We assign their metallicities on a scale of m to m– – – – –, where m is indistinguishable from the main sequence standard and 6 “-” indicate the most severely metal poor subdwarf.

As metallicity drops, the TiO5 band gradually weakens, as predicted in the model spectra of Figure 9 for stars cooler than  $\sim 4000\text{K}$ . Our M1.0V standard (metallicity m), LHS 109 (metallicity scale m– – – – –), and LHS 518 (metallicity scale m– – – – –) have trigonometric parallaxes, and as predicted from the models, their positions on the HR diagram shift bluer with decreasing metallicity. At metallicity m–, SCR 2101–5437 is slightly metal poor compared to M1.0V (TiO5), and has slightly higher gravity than M1.0V (CaH). Unfortunately, none of the SCR objects in Figure 19 have trigonometric parallaxes, yet. Thus, they cannot be plotted on the HR diagram, and the progressive trend in metallicity effects for these stars cannot yet be shown.

*Gravity effects* The effects of gravity can be seen at three different metallicities for type M1.0VI. Spectra for metallicity scale m– – – – – subdwarfs are shown in the middle of Figure 19. SCR 0701–0655 has higher gravity than LHS 518. Both spectra are identical, except at CaH. Spectra for metallicity scale m– – – – – subdwarfs are shown in the bottom of Figure 19. A clear trend can be seen. Three of these five objects have trigonometric parallaxes — LHS 109, LHS 385 and LHS 335 — and their gravity differences shift their locations on the HR diagram toward redder and less luminous territory, as with previously discussed types. (LHS 385’s high parallax error is the likely cause of the slight offset between its HR diagram position and LHS 109’s.) Finally, as shown at the top of Figure 20 for metallicity scale m– – –, SCR 0654–7358 has higher gravity than SCR 1756–5927.

**LHS 440** was reported by Bidelman (1985) to be type M1.0V. The second set of spectra in Figure 20 shows that LHS 440 has deeper absorptions in all three CaH bands than the M1.0V standard, but is virtually identical at all other wavelengths. This indicates that LHS 440 probably has higher gravity than M1.0V, making it less luminous than main sequence stars on the HR diagram. Because it is just barely one magnitude below the main sequence fit, we currently assign it a type of M1.0VI:.

**LHS 1970 and LHS 2734B** both have noisy spectra because they are faint. As shown in Figure 20, their strong CaH bands indicate that they are high gravity (strong CaH), low metallicity (weak TiO5) subdwarfs. LHS 1970’s location on the HR diagram confirms that it is a subdwarf and Gizis (1997) reported its spectral type is esdM2.5. LHS 2734B’s spectral similarity to LHS 1970 indicates that it also has low metallicity and high

gravity. Unfortunately, we measure zero parallax for the LHS 2734AB pair, within the errors, so we can not plot them on the HR diagram (see section 7.2.3 for discussion of LHS 2734A). We assign both LHS 1970 and LHS 2734B types of M1.0VI: because of their noisy spectra.

**LHS 158** has a spectrum nearly identical to an M1.0V star, but falls one magnitude below the main sequence line on the HR diagram. It also has  $V_{tan} = 191 \text{ km sec}^{-1}$  (Jao et al. 2005). The low luminosity and tangential velocity together imply that LHS 158 may be a subdwarf, so we assign it a type of M1.0[VI], representing its uncertain assignment as a subdwarf, as is the case for the mid K-type subdwarfs.

### 7.2.7. M2.0VI

This is the earliest type for which we see that CaH1 absorption deepens as the metallicity decreases, as shown by the curves in Figure 8 for  $T_{eff}$  less than 3500K. Figure 21 shows the spectra for stars discussed in this section.

**LHS 406, GJ 191 (Kapteyn’s Star) and WT0233** form a sequence of decreasing metallicity. These three stars have parallaxes, and their positions on the HR diagram show a clear trend toward the blue for stars with decreasing metallicity (shown with open arrows). One of the prototypes of the subdwarf class, Kapteyn’s Star, which has  $[\text{Fe}/\text{H}] = -0.99$  from Woolf & Wallerstein (2005), has previously been reported to be a type sdM1.0 in Gizis (1997). We assign a somewhat later type for this famous star, which at 3.9 pc is the nearest known subdwarf of any type.

**GJ 191/LHS 3620 and LHS 406/LHS 127** The relative effects of gravity can be seen in the spectra of these two pairs. LHS 3620 has much stronger gravity than GJ 191, while LHS 127 has slightly stronger gravity than LHS 406. On the HR diagram, LHS 3620 is redder and less luminous than GJ 191, as expected. LHS 162 has a spectrum (not plotted) virtually identical to LHS 3620, but is only slightly redder and less luminous than GJ 191.

**LHS 318** is almost identical to GJ 191 except for weaker TiO5 absorption. This weaker TiO5 indicates LHS 318 is slightly more metal poor than GJ 191, so the blue end of the spectrum should be “brightened”; however, it is not. LHS 318’s location on the HR diagram in relation to GJ 191 suggests a higher gravity that is not apparent in its spectrum. We would expect LHS 318 to appear brighter and bluer than GJ 191 if metallicity is the only difference in these two stars, so we assign it as M2.0VI.

### 7.2.8. *M3.0VI*

**LHS 228/LHS 189AB and LHS 326/LHS 398** are pairs of stars with nearly identical spectra. As shown in Figure 22, LHS 326, LHS 228 and our M3.0V standard form a metallicity sequence. As in the case of M2.0VI, CaH1 absorption deepens for M3.0VI stars with decreasing metallicity. In the case of LHS189AB, Costa et al. (2006) reported their separation is about  $3''$  and their  $\Delta R \sim 0.5\text{mag}$ . At a distance of 22.1 pc, the projected separation implies a distance of  $\sim 66$  AU between the two components. This spectrum has combined both components, so we assign it type M3.0VI: until completely “clean” data can be acquired for the components, and their individual metallicity scales can be determined.

**WT 135** was previously identified by Henry et al. (2002) as type M2.5V. We assign its metallicity m–: because its TiO5 band is the same as seen in LHS 228, but the blue end of its spectrum is “brightened” (making it somewhat metal poor compared to LHS 228). It lacks a trigonometric parallax, so we cannot plot it on the HR diagram.

**LHS 272** with  $V - I = 4.29$  is similar in color to WT 135 ( $V - I = 4.32$ ) but appears to have slightly higher gravity (deeper CaH bands). Overall, the blue end of its spectrum (6000Å–6300Å) is comparable to WT 135, although with portions slightly brighter and portions slightly fainter. We therefore assign its metallicity to be the same as WT 135, at m–:. It is slightly redder and less luminous than LHS 228. We use both hollow (metallicity) and solid (gravity) arrows to indicate its location relative to LHS 228. The competition between metallicity and gravity effects in this case seems to indicate that gravity is the dominant factor because LHS 272’s position moves to the red, rather than the blue. LHS 272 was reported as sdM3.0 in Gizis (1997).

**LHS 326, SCR 2204–3347/LHS 541 and SCR 1916–3638** form a clear sequence of gravity effects. Other than having a noisier spectrum (not shown), LHS 541 appears to be identical to SCR 2204–3347. LHS 541’s location on the HR diagram relative to LHS 326 reflects the effects of gravity.

### 7.2.9. *M3.5VI*

This half type is assigned because four stars have spectra shown in Figure 23 with redder slopes than M3.0VI (compare to LHS 228, a M3.0VI subdwarf in the second set of spectra), but not as steep as M4.0VI (see next section). Conveniently, all four stars of this type have trigonometric parallaxes.



**LHS 381 and LHS 144** have lower metallicities than observed for our M3.5V standard. Their spectra redward of 8200Å match M3.5V relatively well and show the same trends predicted by the GAIA models (Figure 9) for stars with temperatures of about 3200K. Their CaH1 indices also match the trend discussed in section 4.1.

**LHS 375 and LHS 515** have higher gravities than LHS 381 and LHS 144, respectively. Both objects are redder and less luminous than their low gravity counterparts on the HR diagram. Gizis (1997) reported LHS 375 to be an esdM4.0 subdwarf and Reid and Gizis (2005) reported LHS 515 to be an esdM5.0 subdwarf. We classify both as M3.5VI with high gravities.

#### 7.2.10. *M4.0VI to M6.0VI*

We identify only four subdwarfs with types M4.0VI to M6.0VI, only one of which, LHS 2067A, currently has a trigonometric parallax. Because of the paucity of such objects, we do not yet have a sample sufficiently large to map out the effects of metallicity and gravity.

When stars are cooler than  $\sim 3200\text{K}$ , their spectra redward of 7500Å change radically between  $[\text{m}/\text{H}]=0.0$  and  $-1.0$  (see Figure 9), but are rather more stable between  $[\text{m}/\text{H}] = -1.0$  and  $-2.0$ . For example, Figure 24 shows comparisons between LEHPM 3861 (M4.0VI) and our M3.0V and M4.0V standards. Clearly, if its spectral type is incorrectly assigned to be M3.0VI, there is excess red flux. Lodieu et al. (2005) reported LEHPM 3861 to be a sdM6.0 subdwarf.

**LEHPM 3861** (M4.0VI), **LHS 1490** (M5.0VI), **LHS 334** (M6.0VI), and **LHS 2067A** (M6.0VI), whose spectra are shown in Figure 24 are assigned subdwarf spectral types later than M3.5VI. The types are based on fluxes redward of 7500Å, with emphasis on three pseudo-continuum peak points at 8130, 8250 and 8840Å (marked in Figure 24 with dotted lines). As shown in Figure 9 for stars cooler than 3200K, the flux decreases at these three points as the metallicity drops from  $[\text{m}/\text{H}]=0.0$  to  $-1.0$  and this trend becomes even more prominent as  $T_{\text{eff}}$  decreases. Thus, these three peaks for a low metallicity subdwarf will not be brighter than a dwarf with a comparable spectral type.

**LHS 2067A** is bluer than M6.0V on the HR diagram and is slightly below the main sequence line. LHS 2067A was previously identified to be a subdwarf in Kirkpatrick et al. (1995), but no spectral type was given. Although its CaH1 and CaH2+CaH3 indices are not in the subdwarf region (see Figure 3 and Table 1), its spectrum is clearly different from M6.0V and M7.0V, with stronger CaH1 absorption. It forms a wide ( $\sim 55''$  NE) common proper motion pair with a white dwarf, LHS 2067B, so it, like LHS 193AB and LHS 300AB,

forms an intriguing pair that can be used for comparing metallicity and white dwarf ages. At the system’s distance of 25.5 pc, the large separation of the pair corresponds to 472 AU, indicating that significant pollution of the subdwarf by the white dwarf during its planetary nebula phase seems unlikely.

**LHS 334** was reported to be a sdM4.5 subdwarf in Reid and Gizis (2005). We assign it a type of M6.0VI because of the good match of the three pseudo-continuum peak points to LHS 2067A, which appears to be slightly more metal rich than LHS 334.

## 8. Application to SDSS Subdwarfs

It is useful to apply our spectral typing methodology to the recent work of West et al. (2004), who have provided a significant sample of sixty new subdwarf spectra, all acquired and reduced in a homogeneous way. The subdwarfs were selected and spectral types were assigned via their CaH and TiO indices. For comparison, we retrieved fifty<sup>8</sup> of the publicly available spectra and use our new method to assign types. The brightest star among these subdwarfs has  $r=17.0$ , resulting in somewhat noisy spectra for the sample, so we smoothed the spectra by averaging the flux over five pixels and normalizing at 7500Å. Representative spectra are shown in Figures 25 and 26.

The SDSS spectra have telluric lines removed, so we use M dwarf standard spectra from Bochanski et al. (2007) that also omit the telluric lines, rather than our own standards. In addition, Bochanski et al. (2007) do not present half-type standard spectra, so we do not present any half-type spectra for the SDSS subdwarfs. Previous and current types are listed in Table 3. The subdwarfs have types between M1.0 and M3.0 in Bochanski et al. (2007), while our efforts yield types between M2.0 and M5.0.

Our point here is not to assign definitive spectral types, but to illustrate the gravity and metallicity effects in the SDSS dataset, and provide examples of the application of our proposed spectral typing method. Stars of a given spectral type, of course, may have a range of metallicities, as discussed at length in previous sections, and as is evident in Figure 25. Gravity effects are also seen, as shown in Figure 26. Three stars (SDSSJ085843.89+511210.1, SDSSJ093141.85+453914.5 and SDSSJ145447.32+011006.8) previously identified as subdwarfs have spectra identical to SDSS M dwarfs. Two other stars, SDSSJ083217.77+522408.2 and SDSSJ113501.76+033720.3, ap-

---

<sup>8</sup>The remaining 10 subdwarfs could not be retrieved using the SDSS DR4 website, even though we used a 1' search radius and coordinates from their table.

pear to have slightly higher gravities (stronger CaH1 line) than dwarfs, but the rest of these two spectra are almost identical to SDSS M dwarfs. Therefore, to these two stars we assign types of M2.0VI: and M3.0VI: with g+., respectively.

Unfortunately, the SDSS dwarfs do not have parallaxes that can be used to confirm their locations on the HR diagram. In addition, we find many SDSS subdwarfs with identical spectral types that have  $g - z$  colors as different as 0.53. It is therefore difficult to verify how metallicities and gravities affect the SDSS absolute magnitudes and colors. Nonetheless, the robust sample of West et al. (2004) provides many additional subdwarfs that can be targeted for further work.

## 9. Discussion

### 9.1. Why Subdwarfs’ Physical Parameters Are Not Listed Explicitly

Stellar spectra follow trends primarily defined by temperature, yet as we have seen, metallicities and gravities also have significant effects on the spectra of cool subdwarfs. Here we have provided a consistent spectral sequence for subdwarfs that is based on linking observed subdwarf spectra to spectra acquired for main sequence dwarfs using the same telescope/instrument/observing protocols. Here we compare synthetic to observed spectra to evaluate how well we can assign values for the temperatures, metallicities, and gravities of cool dwarfs and subdwarfs.

#### 9.1.1. Matching Synthetic and Observed Spectra

To test the reliability of the model grids and our fitting procedures, we use our six standard spectral sequence dwarfs with types M0.0V to M5.0V as test spectra (see Figure 27). All are currently believed to be uncorrupted single red dwarfs because they show no evidence of multiplicity from combinations of (1) HST/NICMOS observations, (2) optical speckle observations, (3) optical CCD imaging, and/or (4) three or more years of astrometric observations that would reveal perturbations from unseen companions that contributed significant light to the system.

We compared the spectral region from 6000Å to 9000Å after normalizing both the GAIA grid spectra and ours at 7500Å, and applying a Gaussian function to the much higher resolution GAIA model spectra to match our resolution. If a star had a significant spectral shift because of radial velocity, we manually offset the spectrum to match features obvious

in the synthetic spectra. We then calculated the reduced  $\chi^2$  differences (hereafter, simply  $\chi^2$ ) between the model grid spectra and ours. Because the synthetic spectra do not have the telluric lines of  $\text{O}_2\alpha$  (6270Å–6330Å),  $\text{O}_2\text{B}$  (6860Å–6980Å),  $\text{O}_2\text{A}$  (7590Å–7710Å) and water (7150Å–7330Å and 8952Å–9000Å), these absorption regions were excluded in the  $\chi^2$  calculations.

The selected model grids have effective temperatures of 2400K to 4500K,  $[\text{m}/\text{H}]$  from  $-2.0$  to  $+0.5$ , and  $\log g$  from 4.0 to 5.5 in steps of 100K, 0.5 dex, and 0.5 dex, respectively. The upper panel of Figure 10 shows that for a metallicity of  $-1.0$  and temperature 3500K, varying the gravity causes changes in only certain wavelength regions (and some specific lines). This means  $\chi^2$  is only sensitive to  $\log g$  in relatively small spectral regions, but not for the overall spectrum. On the other hand, as shown in the lower panel of Figure 10, for the same temperature star with  $\log g$  fixed at 5.0, there are large differences between spectra when the metallicity is varied.

Thus, for a star of a given temperature, the gravity changes the overall spectral shape minimally while the metallicity changes it a great deal, so we first secure a star’s metallicity and then its gravity. Figure 28 is an example for M1.0V standard star, showing  $\chi^2$  curves at various metallicities and gravities. Each curve represents a specific  $\log g$  and grid spectra have temperatures incremented by 100K. The smallest scatter in the plots is found when  $[\text{m}/\text{H}]$  is  $-0.5$ , which is adopted as the star’s metallicity. We then examine each point in the  $[\text{m}/\text{H}]=-0.5$  panel to find the best fit with the smallest  $\chi^2$  at this fixed metallicity. In this case, the best fit has  $T_{\text{eff}} = 3600\text{K}$ ,  $\log g = 4.5$  and  $[\text{m}/\text{H}] = -0.5$ .

### 9.1.2. Discrepancies Between the Best Fitting Synthetic and Observed Spectra

In this paper we do not explicitly list temperatures, metallicities, or  $\log g$  values for subdwarfs. A few examples support our reasoning for not listing these physical parameters. Figure 27 shows observed spectra and the best fit synthetic spectra for six main sequence spectral standard stars with types M0.0V to M5.0V (in steps of 1.0 subtypes). In general, the overall slopes of the model spectra fit fairly well, especially for the earlier types. However, we discuss here six regions labeled at the top of Figure 27 that do not match, which are particularly relevant for the subdwarfs that are the focus of this paper.

1. Region 1: The observed spectrum is always less luminous than the model, except for M0.0V.
2. Region 2: This is the CaH1 absorption region. Observed spectra have shallower CaH1 features than the models. The depth of CaH1 is determined by a combination of gravity

and metallicity. Decreasing CaH1 absorption can be caused by either decreasing gravity or decreasing metallicity. However, fine tuning the gravity or metallicity affects not only the CaH1 feature, but the TiO5 feature in region 4.

3. Region 3: The pseudo-continuum always peaks near 6530Å in the observed spectra. However, the synthetic spectra have this peak “red-shifted” to ~6650Å.
4. Region 4: Containing the CaH2, CaH3 and TiO5 features, this is the most important region for examining the interplay of metallicity and gravity in cool dwarfs. Unfortunately, the CaH2 feature is blended with O<sub>2</sub>B so it is not entirely reliable for analysis of spectra taken through the Earth’s atmosphere. The CaH3 and TiO5 features are usually weaker in observed spectra than in the models. The strength of TiO5 is primarily driven by metallicity, not gravity (see the bottom panel of Figure 10). Therefore, this region reveals valuable information about a star’s metallicity, in particular at types later than M2.0.
5. Region 5: Several Fe I absorption lines (8388Å, 8440Å and 8718Å) and a Mg I line (8718Å) in the model spectra redward of 8300Å are deeper than observed. This indicates poor metallicity matches and/or poor modeling of those particular lines.
6. Region 6: Overall, redward of 8000Å the M4.0V and M5.0V matches are not as good as other regions. This likely indicates fundamental problems with the strengths of some opacity sources (e.g. H<sub>2</sub>O and TiO) in the models.

The top panel of Figure 29 shows the two best fitting synthetic spectra for our M1.0V standard. The red spectrum (3600K,  $[m/H] = -0.5$ ,  $\log g = 4.5$ ) provides the best fit and the blue spectrum (3400K,  $[m/H] = -1.0$ ,  $\log g = 4.0$ ) is second best. There are only slight differences between the two synthetic spectra — the CaH1 strengths and continuum fluxes redward of 8000Å are the only notable differences. However, the effective temperatures for the two models differ by 200K, while the metallicities and  $\log g$  each differ by 0.5 dex. The lower panel of Figure 29 shows the two best fit spectra for our M3.0V standard. The best fit (red line) yields 3200K,  $[m/H] = -0.5$ ,  $\log g = 4.5$  and the second best fit (blue line) yields 3300K,  $[m/H] = 0.5$ ,  $\log g = 5.5$ . However, neither model fit is an ideal match to the observed spectrum (which is why the  $[m/H]$  value for the two best matches differs by an order of magnitude), and matches become even poorer for cooler stars.

As a whole, many of the best fits for the main sequence dwarfs in Figure 27 are for metallicities of  $-0.5$ , which is somewhat lower than studies that have specifically attempted to assign metallicities to cool dwarfs in the solar neighborhood. For example, the mean  $[Fe/H]$  from 21 M dwarf secondaries in Bonfils et al. (2005) is  $-0.09$ , and the mean  $[Fe/H]$

from five M dwarf secondaries in Bean et al. (2006) is  $-0.17$ . Also worthy of note is that the fits were made for metallicities incremented by 0.5 dex, and stars with metallicities between  $-0.5$  and  $0.0$  may have slightly better fits for  $-0.5$ . One might think that interpolation from existing model grids would allow better fits. There are, however, degeneracy problems in matching model spectra to observations — when fitting the three-dimensional space of temperature, metallicity, and gravity, two or more synthetic spectra yield low points with similar  $\chi^2$ , as is the case for the M1.0V example discussed above. More worrisome is that the overall discrepancies discussed above (items 1 to 6) cannot be removed through interpolation in the existing model grids.

Figure 30 illustrates how the  $\chi^2$  values change for best fit matches of the GAIA models to our spectral standards with types M0.0V through M5.5V. Spectral type M3.0V is the latest type for which a reasonably tight plot like the one shown in Figure 28 can be identified. The dotted line in Figure 30 provides a dividing point between reasonable matches and poor matches, indicating that once  $\chi^2$  exceeds 10, the  $\chi^2$  plots are too scattered to choose a unique set of model spectra parameters to match observed spectra.

A specific example of applying the models to one of our subdwarfs illustrates the large discrepancies that must be overcome to derive reliable parameter values for metallicity and gravity. When we applied our fitting algorithm to our observed spectrum for LHS 335 (M1.0VI, shown in Figure 31), we discovered that the “best fit” was quite poor. Although the formal  $\chi^2$  value was 5.9 (less than our cutoff of 10, indicating a reliable fit), the model’s CaH3 band is not deep enough, the TiO5 band is too deep, and the continuum flux redward of  $7700\text{\AA}$  is less than observed.

For myriad reasons we conclude that we cannot strictly determine reliable metallicities and  $\log g$  values for cool dwarfs using the fitting method discussed here. Thus, until improved model grids are available, we defer assignment of numerical values for temperatures, metallicities and  $\log g$  values. However, we certainly *can* use the GAIA model grids to mimic *trends* (discussed in §5 and 6) in the spectra of subdwarfs to compare the stars within a framework of changing physical parameters.

## 9.2. The Confusion Between *sd*, *esd* and *usd* Prefixes

Gizis (1997) proposed that subdwarfs have two subclasses, “subdwarfs” (*sd*) and “extreme subdwarfs” (*esd*) based on their CaH and TiO5 band strengths. He separated the two classes using the line shown in the top panel of Figure 3. Another term, “ultra subdwarf”, was proposed by Caldwell et al. (1984) to describe GJ 59B ( $V - K_s = 3.42$ ,  $M_{K_s} = 7.89$ ),

which is  $\sim 2.5$  magnitudes underluminous compared to main sequence stars of similar color. This star falls in the region including “extreme subdwarfs” in Gizis (1997). More recently, Lépine et al. (2007) adopted the “ultra subdwarf” (*usd*) term for subdwarfs found to have stronger CaH features than “extreme subdwarfs”. We believe these terms confuse the situation and do not address the underlying astrophysics. We recommend that they not be used for the following reasons.

- These terms can only be applied to M-type subdwarfs, not K-type subdwarfs. Both observed spectra and models show that K subdwarfs do not follow the trends in CaH absorption with metallicity that M subdwarfs follow. Thus, there is no clear delineation for K-type subdwarfs.
- Empirically, the values of the indices are affected by a complicated interplay of temperature, metallicity, and gravity effects. One cannot separate these three factors simply based on the indices. Typically, when the term “extreme” subdwarf is used, it refers to “very low metallicity” alone. But that is only one part of the portrait that needs to be painted for a given subdwarf. For example, filled boxes shown in Figure 4 indicate stars previously identified as extreme subdwarfs from their indices. Note that the spread for “extreme” subdwarfs at a given color ( $V - K_s \sim 3.5$ ) is 3.5 magnitudes, or a factor of  $\sim 25$  in luminosity. There is no clear separation in the fundamental HR diagram between stars termed “subdwarfs” and “extreme subdwarfs” classified using spectral indices.

In Figure 32, we take a detailed look at astrophysical causes that shift points in the CaH2+CaH3/TiO5 indices plot. Red circles in Figure 32 represent the lowest metallicity stars of each spectral type we have presented here. Some of these very low metallicity stars are not located in the “extreme” subdwarf region. Green circles represent subdwarfs in our sample that have highest gravity at a given type. Solid lines connect stars having the same metallicity rankings in Table 2 (components of the common proper motion binary LHS 2734 AB are connected by a dotted line because they presumably have the same metallicity). It is clear from these three pairs that high gravity can push a subdwarf toward or into the extreme subdwarf region, even if the metallicities of the two objects are similar. In addition, many other high gravity subdwarfs (green circles) that do not have particularly low metallicities are also in the subdwarf region. Finally, the two blue circles represent LHS 440 and SCR 1822-45542, which subdwarfs having the same metallicities but higher gravities than dwarfs. Their spectroscopic features indicate that they are subdwarfs, but they do not have low metallicity at all. Yet, they fall in the subdwarf region of the indices plot, which has traditionally indicated low metallicity.

The inset in Figure 32 helps explain this phenomenon. We have calculated the spectral indices from the GAIA synthetic spectra with  $[m/H]=0.0$ ,  $-1.0$  and  $-2.0$  and temperatures of 2800–4400K. Two different gravities were selected for each metallicity. The models indicate that both low metallicity and high gravity push stars toward the extreme subdwarf region. Thus, if two stars have the same metallicity, the one with higher gravity will be pushed more toward the extreme subdwarf region, indicating that this spectral indices plot is not a clear indicator of metallicity alone.

- Theoretically, in order to show how complicated the TiO5 vs CaH1 plot is astrophysically, we use GAIA model grids to do a demonstration. We have chosen models with  $2700\text{K} < T_{eff} < 4500\text{K}$ ,  $-3.0 < [m/H] < 0.0$  and  $4.0 < \log g < 5.5$ , and have calculated the output CaH and TiO5 indices, as shown in Figure 33. As is apparent in the plots, for a given (TiO5, CaH1) indices pair, there are many possible parameter combinations. Both indices are a function of temperature, gravity and metallicity, which would require a 6-D plot to describe TiO5 vs CaH. Unfortunately, one cannot classify stars as subdwarfs, extreme subdwarfs, or ultra-subdwarfs using 2-D index plots that infer trends in metallicity alone.

For these reasons, the indices only indicate a star’s location on these particular spectroscopic indices plots. They do not address low or very low metallicity only, as implied by past usage, but instead incorporate effects of both metallicity and gravity for stars of a given temperature. They do not provide obvious or direct links to positions on HR diagrams, which provide the astrophysical meaning underlying stellar classification. Finally, the modifiers “extreme” and “ultra” themselves have effectively the same meaning and do not provide meaningful information about their differences.

### 9.3. Why Previous Methods Work for Dwarfs but Not Subdwarfs

In the Palomar-MSU spectroscopy survey, Reid et al. (1995) used the TiO5 index and polynomial equations to assign subtypes for M dwarfs. Gizis (1997) then applied the same methodology to M subdwarfs. As can be seen in Figure 33, TiO5 versus temperature is effectively linear, especially for  $\log g = 5.0$  and  $[m/H] = 0.0$ , so this typing method works successfully for a fairly homogeneous set of dwarfs (possibly with somewhat different gravities) — temperature is the main factor affecting the overall slope of red dwarf spectra. However, it is not as simple for subdwarfs because both low metallicity and high gravity stars are sorted into the same TiO5 values for various combinations. Thus, the same methodology that works for dwarfs cannot straightforwardly be applied to subdwarfs.



#### 9.4. The Subdwarf Spectral Standards from Lépine et al. (2007)

Lépine et al. (2007) recently released a set of spectral standards for the sdM, esdM and usdM subclasses using wider spectral coverage than used by Gizis (1997) and revised polynomial equations. Their project is contemporaneous with ours, but unfortunately there is only one star in both samples, LHS 228. We assign it a type of M3.0VI while their type is sdM2.0. As shown in Figure 34, gravity effects come into play in their spectra<sup>9</sup>. Each pair of spectra shown have nearly identical continua, with CaH bands being the only significant spectral difference. Using their method, each of the pairs of subdwarfs shown is assigned a different sub-type. We believe that by using the overall shape of the spectrum, with knowledge that both metallicity and gravity affect certain regions, each pair should have the same spectral sub-types. Based on the trends from GAIA models, the pairs of spectra shown have different gravities, as we have also seen in our spectra and those from SDSS (shown in §8). Figure 35 illustrates spectral differences at types M3.0, M5.0 and M7.5. Although the spectra in each panel have the same sub-types for sdM, esdM and usdM subdwarfs, they show very different continua between 6000Å and 9000Å. When examining the overall spectra, there is no clear morphology trend for each sub-type. We conclude that using spectral indices alone omits important information evident in the overall morphology of the spectra that is useful to spectral classification.

### 10. Conclusions

We have discussed 88 cool subdwarfs using spectra covering 6000–9000Å. Based on these spectra and the trends from GAIA model grids, we have redefined the subdwarf spectral sequence, spanning types K3.0 to M6.0. We find that wide spectral coverage is the key to defining a subdwarf’s spectral type. We consider this to be an important, but not final, step in defining the subdwarf spectral sequence.

Through the understanding of GAIA model grids, we find that the key to assigning a subdwarf spectral type is to compare the spectrum to dwarf spectral standards in regions affected minimally by metallicity and gravity, thereby making a direct link between the dwarf and subdwarf sequences. Even so, it remains difficult to establish a definitive sequence for subdwarfs because of the multi-faceted nature of their spectra. Until we have surveyed a large number of subdwarfs and covered a multitude of possible temperatures, metallicities, and gravities, a definitive sequence will remain elusive.

---

<sup>9</sup>Their telluric lines have been removed.

From an analysis of the history of the term “subdwarf,” and the layout of the fundamental HR diagram, we propose that the suffix “VI” be used, rather than the “sd” prefix, as the preferred spectral classification notation. This reduces the confusion between cool subdwarfs and hot OB subdwarfs. This is also prudent because subdwarfs really do form an independent class of stars on the HR diagram, for which five Roman numerals are in common use, with cool subdwarfs naturally falling beneath the main sequence V types.

Overall, we find that trigonometric parallaxes are crucial for identifying mid K-type subdwarfs, and allow us to understand how the complex interplay of temperatures, metallicities, and gravities affects the positions of cool dwarfs on the HR diagram. We found that mid K-type subdwarfs can not be identified spectroscopically using our data. There are many G and M-type subdwarfs, so K-type subdwarfs presumably exist, but additional information is needed to identify them. Consequently, we use metallicities, kinematics, and parallaxes to ascertain their true natures and use their locations on the HR diagram to flag them as “subdwarfs” (other spectroscopic wavelength and resolution combinations could also be used). Because they are not spectroscopically identified as subdwarfs in our spectra, we use a conservative notation, [VI], for subdwarfs of types K3.0 through K5.0 to indicate their questionable status.

We have confirmed that spectroscopic indices are useful in separating late K to late M-type subdwarfs from dwarfs, but that the indices have limitations when attempting to understand the astrophysical causes leading to observed subdwarf spectra. When combined with trigonometric parallax and photometric information, our results show that for lower metallicities, subdwarfs are generally bluer and brighter at optical wavelengths, so they slide up and to the left on the HR diagram (using axes  $V - K_s$  vs  $M_{K_s}$ ). In contrast, higher gravities make stars redder and less luminous at optical wavelengths, so subdwarfs generally slide down and to the right on the same HR diagram. Because of the complex, and not yet completely mapped out, interplay of temperatures, metallicities, and gravities, we conclude that the “extreme” and “ultra” prefixes *only* outline locations on spectroscopic indices plots, and do not successfully differentiate the underlying astrophysical causes for shifts on CaH/TiO5 plots.

Improvements in the subdwarf spectral sequence can be made by observing wide common proper motion subdwarf binaries, like LHS 2734AB discussed here. Assuming identical metallicities, such binaries allow us to constrain one of the three variables that affects cool dwarf spectral types. The ultimate subdwarf spectral sequence will be three-dimensional, with temperature, metallicity, and gravity as independent variables (see Kirkpatrick 2005, Figure 11).

Previously, gravity effects in cool subdwarfs have been almost entirely ignored, as metal-

licity was the factor considered to describe changing CaH features. Our results clearly show the importance of gravity effects. An additional factor, contamination from an unseen, evolved, i.e. white dwarf, companion could also change the slope of subdwarf spectra. Thus, comprehensive surveys of subdwarfs for companions are warranted<sup>10</sup>. Finally, this work also shows that current synthetic spectra provide a useful framework in which to evaluate cool dwarf spectra, but they do not yet provide perfect matches, indicating that atmospheric models still require fine-tuning to make additional advances in the future characterization of cool dwarfs (P. Hauschildt private communication).

We would like to thank the referee, Sandy Leggett, for her very helpful comments that improved this paper. We also thank Pat Boeshaar and Doug Gies for their comments and suggestions. Peter Hauschildt was instrumental in helping us use the GAIA model grids, and Vincent Woolf contributed the old version of synthetic spectra. We thank Sebastien Lepine for providing his subdwarf spectra to us before they were publicly available, and for his comments on section 9.4. We appreciate the help of many members of the Georgia State University (GSU) RECONS team in their data acquisition and reduction efforts, especially Charlie Finch and Jennifer Winters. This work has been supported at GSU by NASA’s Space Interferometry Mission, the National Science Foundation (NSF, grant AST-0507711), and GSU. Finally, we wish to thank the other members of the SMARTS Consortium, without whom the telescopes at CTIO used for this effort might not be available.

This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This paper has also made use of data from the SDSS. Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, NSF, the U.S. Department of Energy, NASA, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. This work has used data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center at California Institute of Technology funded by NASA and NSF.

## REFERENCES

Adams, W. S., & Joy, A. H. 1922, *ApJ*, 56, 242

---

<sup>10</sup>We have an ongoing project to survey cool subdwarfs for companions. To date, we have found that cool subdwarfs have a lower binary fraction than cool dwarfs as reported by Riaz et al. (2008), so the contamination assumption from possible unseen companions appears unlikely.

- Adams, W. S., Joy, A. H., Humason, M. L., & Brayton, A. M. 1935, *ApJ*, 81, 187
- Allard, F., & Hauschildt, P. H. 1995, *ApJ*, 445, 433
- Allen, C., Poveda, A., & Herrera, M. A. 2000, *A&A*, 356, 529
- Bean, J. L., Sneden, C., Hauschildt, P. H., Johns-Krull, C. M., & Benedict, G. F. 2006, *ApJ*, 652, 1604
- Beaulieu, T. D., in preparation, 2009
- Bessell, M. S. 1982, *Proceedings of the Astronomical Society of Australia*, 4, 417
- Bessel, M. S. 1990, *A&AS*, 83, 357
- Bidelman, W. P. 1985, *ApJS*, 59, 197
- Bochanski, J. J., West, A. A., Hawley, S. L., & Covey, K. R. 2007, *AJ*, 133, 531
- Boeshaar, P. C. 1976, Ph.D. Thesis
- Bonfils, X., Delfosse, X., Udry, S., Santos, N. C., Forveille, T., & Ségransan, D. 2005, *A&A*, 442, 635
- Brott, I., & Hauschildt, P. H. 2005, *The Three-Dimensional Universe with Gaia*, 576, 565
- Burgasser, A. J., et al. 2003, *ApJ*, 592, 1186
- Burgasser, A. J., & Kirkpatrick, J. D. 2006, *ApJ*, 645, 1485
- Caldwell, J. A. R., Spencer Jones, J. H., & Menzies, J. W. 1984, *MNRAS*, 209, 51
- Carney, B. W., Latham, D. W., Laird, J. B., & Aguilar, L. A. 1994, *AJ*, 107, 2240
- Cayrel de Strobel, G., Soubiran, C., & Ralite, N. 2001, *A&A*, 373, 159
- Cayrel de Strobel, G., Soubiran, C., Friel, E. D., Ralite, N., & Francois, P. 1997, *A&AS*, 124, 299
- Chamberlain, J. W., & Aller, L. H. 1951, *ApJ*, 114, 52
- Costa, E., & Méndez, R. A. 2003, *A&A*, 402, 541
- Costa, E., Méndez, R. A., Jao, W.-C., Henry, T. J., Subasavage, J. P., Brown, M. A., Ianna, P. A., & Bartlett, J. 2005, *AJ*, 130, 337

- Costa, E., Méndez, R. A., Jao, W.-C., Henry, T. J., Subasavage, J. P., & Ianna, P. A. 2006, *AJ*, 132, 1234
- Cox, N. 2000, *Allen’s astrophysical quantities*, New York, AIP Press, Springer
- Deacon, N. R., Hambly, N. C., & Cooke, J. A. 2005, *A&A*, 435, 363
- Digby, A. P., Hambly, N. C., Cooke, J. A., Reid, I. N., & Cannon, R. D. 2003, *MNRAS*, 344, 583
- Gizis, J. E. 1997, *AJ*, 113, 806
- Gizis, J., & Reid, I. 1997, *PASP*, 109, 1233
- Gizis, J. E., & Reid, I. N. 1999, *AJ*, 117, 508
- Gray, R. O., Corbally, C. J., Garrison, R. F., McFadden, M. T., & Robinson, P. E. 2003, *AJ*, 126, 2048
- Gray, R. O., Corbally, C. J., Garrison, R. F., McFadden, M. T., Bubar, E. J., McGahee, C. E., O’Donoghue, A. A., & Knox, E. R. 2006, *AJ*, 132, 161
- Greenstein, J. L., & Eggen, O. J. 1966, *Vistas in Astronomy*, 8, 63
- The *Hipparcos* and Tycho Catalogues, 1997, ESA SP-1200 (Noordwijk: ESA)
- Feige, J. 1958, *ApJ*, 128, 267
- Giclas, H. L., Burnham, R., & Thomas, N. G. 1971, Flagstaff, Arizona: Lowell Observatory, 1971
- Giclas, H. L., Burnham, R., Jr., & Thomas, N. G. 1978, *Lowell Observatory Bulletin*, 8, 89
- Greenstein, J. L. 1966, *ApJ*, 144, 496
- Greenstein, J. L., & Münch, G. 1966, *ApJ*, 146, 618
- Goldberg, D., Mazeh, T., Latham, D. W., Stefanik, R. P., Carney, B. W., & Laird, J. B. 2002, *AJ*, 124, 1132
- Hartwick, F. D. A., Cowley, A. P., & Mould, J. R. 1984, *ApJ*, 286, 269
- Hawley, S. L., Gizis, J. E., & Reid, I. N. 1996, *AJ*, 112, 2799
- Henry, T. J., Walkowicz, L. M., Barto, T. C., & Golimowski, D. A. 2002, *AJ*, 123, 2002

- Henry, T. J., Subasavage, J. P., Brown, M. A., Beaulieu, T. D., Jao, W.-C., & Hambly, N. C. 2004, *AJ*, 128, 2460
- Henry, T. J., Jao, W.-C., Subasavage, J. P., Beaulieu, T. D., Ianna, P. A., Costa, E., & Méndez, R. A. 2006, *AJ*, 132, 2360
- Heber, U. 1992, *LNP Vol. 401: The Atmospheres of Early-Type Stars*, 401, 233
- Houk, N., & Cowley, A. P. 1975, *Ann Arbor: University of Michigan, Department of Astronomy*, 1975
- Humason, M. L., & Zwicky, F. 1947, *ApJ*, 105, 85
- Jacoby, G. H., Hunter, D. A., & Christian, C. A. 1984, *ApJS*, 56, 257
- Jao, W.-C., Henry, T. J., Subasavage, J. P., Bean, J. L., Costa, E., Ianna, P. A., & Méndez, R. A. 2003, *AJ*, 125, 332
- Jao, Wei-Chun, PhD Dissertation, 2004.
- Jao, Wei-Chun, Henry, T. J., Subasavage, J. P., Brown, M. A., Ianna, P. A., Bartlett, J. L., Costa, E., & Méndez, R. A. 2005, *AJ*, 129, 1954
- Jao, Wei-Chun, 2008, in preparation
- Jaschek, C., & Jaschek, M. 1987, *Cambridge: University Press*, 1987
- Joy, A. H. 1947, *ApJ*, 105, 96
- Kazarovets, E. V., Samus, N. N., Durlevich, O. V., Kireeva, N. N., & Pastukhova, E. N. 2006, *Informational Bulletin on Variable Stars*, 5721, 1
- Kirkpatrick, J. D., Henry, T. J., & McCarthy, D. W., Jr. 1991, *ApJS*, 77, 417
- Kirkpatrick, J. D., Henry, T. J., & Simons, D. A. 1995, *AJ*, 109, 797
- Kirkpatrick, J. D. 2005, *ARA&A*, 43, 195
- Kleinman, S. J., et al. 2004, *ApJ*, 607, 426
- Kuiper, G. P. 1939, *ApJ*, 89, 548
- Kuiper, G. P. 1940, *ApJ*, 91, 269
- Laird, J. B., Carney, B. W., & Latham, D. W. 1988, *AJ*, 95, 1843

- Leggett, S. K. 1992, *ApJS*, 82, 351
- Leggett, S. K., Allard, F., Dahn, C., Hauschildt, P. H., Kerr, T. H., & Rayner, J. 2000, *ApJ*, 535, 965
- Lépine, S., Rich, R. M., & Shara, M. M. 2003, *AJ*, 125, 1598
- Lépine, S., & Shara, M. M. 2005, *AJ*, 129, 1483
- Lépine, S., Rich, R. M., & Shara, M. M. 2007, *ApJ*, 669, 1235
- Lodieu, N., Scholz, R.-D., McCaughrean, M. J., Ibata, R., Irwin, M., & Zinnecker, H. 2005, *A&A*, 440, 1061
- López-Morales, M. 2007, *ApJ*, 660, 732
- Luyten, W. J., *Bruce Proper Motion Survey III. The Stars of Large Proper Motion and the Luminosity Function*, Minneapolis: University of Minnesota, 1939
- Luyten, W. J. 1979, *LHS Catalog*, Minneapolis, University of Minnesota, 1979, 2nd ed.
- Mould, J. R. 1976, *A&A*, 48, 443
- Monet, D. G., Dahn, C. C., Vrba, F. J., Harris, H. C., Pier, J. R., Luginbuhl, C. B., & Ables, H. D. 1992, *AJ*, 103, 638
- Monteiro, H., Jao, W.-C., Henry, T., Subasavage, J., & Beaulieu, T. 2006, *ApJ*, 638, 446
- Morrison, H. L., et al. 2003, *AJ*, 125, 2502
- Nissen, P. E., & Schuster, W. J. 1997, *A&A*, 326, 751
- Nordström, B., et al. 2004, *A&A*, 418, 989
- Öhman, Y. 1934, *ApJ*, 80, 171
- Peterson, R. C., Willmarth, D. W., Carney, B. W., & Chaffee, F. H., Jr. 1980, *ApJ*, 239, 928
- Reid, I. N., van Wyk, F., Marang, F., Roberts, G., Kilkenny, D., & Mahoney, S. 2001, *MNRAS*, 325, 931
- Reid, I. N., Hawley, S. L., & Gizis, J. E. 1995, *AJ*, 110, 1838
- Reid, I. N. and Gizis, J. E., 2005, *PASP*, 117, 676
- Reylé, C., Scholz, R.-D., Schultheis, M., Robin, A. C., & Irwin, M. 2006, *MNRAS*, 373, 705

- Riaz, B., Gizis, J. E., & Samaddar, D. 2008, *ApJ*, 672, 1153
- Rodgers, A. W., & Eggen, O. J. 1974, *PASP*, 86, 742
- Roman, N. G. 1955, *ApJS*, 2, 195
- Ryan, S. G. and Norris, J. E., 1991, *AJ*, 101, 1835
- Sandage, A. R., & Eggen, O. J. 1959, *MNRAS*, 119, 278
- Silva, D. R. and Cornell, M. E., 1992, *ApJS*, 81, 865
- Smart, R. L., Lattanzi, M. G., Jahreiß, H., Bucciarelli, B., & Massone, G. 2007, *A&A*, 464, 787
- Subasavage, J. P., Henry, T. J., Hambly, N. C., Brown, M. A., & Jao, W.-C. 2005, *AJ*, 129, 413
- Subasavage, J. P., Henry, T. J., Hambly, N. C., Brown, M. A., Jao, W.-C., & Finch, C. T. 2005, *AJ*, 130, 1658
- Taylor, B. J., 2001, *A&A*, 398, 731
- Turnshek, D. E., Turnshek, D. A., & Craine, E. R. 1985, *Astronomy and Astrophysics Series*, Tucson: Western Research Company, 1985
- van Altena, W. F., Lee, J. T., & Hoffleit, D. 1995, *The General Catalogue of Trigonometric Stellar Parallaxes* (4th ed.; New Haven: Yale Univ. Obs.)
- Valenti, J. A., Piskunov, N., & Johns-Krull, C. M. 1998, *ApJ*, 498, 851
- Veeder, G. J. 1974, *AJ*, 79, 1056
- Weis, E. W. 1996, *AJ*, 112, 2300
- West, A. A., et al. 2004, *AJ*, 128, 426
- Woolf, V. M., & Wallerstein, G. 2005, *MNRAS*, 356, 963



Table 1. Spectroscopic Indices

| Object       | TiO5  | CaH1  | CaH2  | CaH3  | CaH2+CaH3 |
|--------------|-------|-------|-------|-------|-----------|
| DEN0515–7211 | 1.002 | 1.021 | 1.019 | 1.003 | 2.022     |
| G016–009AB   | 0.973 | 1.018 | 1.021 | 0.998 | 2.019     |
| G022–015     | 0.978 | 0.996 | 0.989 | 0.984 | 1.973     |
| G026–009ACD  | 0.953 | 0.990 | 0.975 | 0.990 | 1.965     |
| GJ0191       | 0.860 | 0.876 | 0.678 | 0.848 | 1.526     |
| GJ0223.1     | 0.958 | 1.001 | 0.981 | 0.971 | 1.952     |
| LEHPM1628    | 0.954 | 0.676 | 0.628 | 0.754 | 1.382     |
| LEHPM3861    | 0.957 | 0.774 | 0.340 | 0.368 | 0.709     |
| LHS0012      | 0.883 | 0.881 | 0.779 | 0.889 | 1.668     |
| LHS0073      | 0.972 | 0.903 | 0.862 | 0.927 | 1.789     |
| LHS0109      | 0.939 | 0.823 | 0.752 | 0.863 | 1.616     |
| LHS0125      | 0.981 | 1.020 | 1.014 | 0.993 | 2.007     |
| LHS0127      | 0.716 | 0.755 | 0.537 | 0.739 | 1.276     |
| LHS0144      | 0.768 | 0.609 | 0.396 | 0.592 | 0.987     |
| LHS0148      | 0.977 | 0.816 | 0.729 | 0.830 | 1.560     |
| LHS0158      | 0.732 | 0.849 | 0.639 | 0.829 | 1.469     |
| LHS0161      | 0.889 | 0.777 | 0.689 | 0.817 | 1.506     |
| LHS0162      | 0.838 | 0.731 | 0.577 | 0.747 | 1.324     |
| LHS0164      | 0.987 | 1.005 | 1.000 | 1.017 | 2.017     |
| LHS0165      | 0.956 | 0.889 | 0.841 | 0.910 | 1.751     |
| LHS0186      | 0.710 | 0.739 | 0.568 | 0.764 | 1.332     |
| LHS0189AB    | 0.630 | 0.733 | 0.492 | 0.719 | 1.211     |
| LHS0193A     | 0.944 | 0.959 | 0.920 | 0.966 | 1.886     |
| LHS0227      | 1.000 | 0.895 | 0.883 | 0.949 | 1.832     |
| LHS0228      | 0.644 | 0.744 | 0.480 | 0.701 | 1.181     |
| LHS0232      | 0.998 | 1.000 | 1.020 | 1.005 | 2.025     |
| LHS0244      | 0.902 | 0.853 | 0.731 | 0.869 | 1.601     |
| LHS0272      | 0.862 | 0.715 | 0.527 | 0.736 | 1.264     |
| LHS0299      | 0.903 | 0.904 | 0.756 | 0.884 | 1.641     |
| LHS0300AB    | 1.031 | 0.933 | 0.886 | 0.943 | 1.830     |
| LHS0318      | 0.934 | 0.747 | 0.617 | 0.797 | 1.413     |

Table 1—Continued

| Object       | TiO5  | CaH1  | CaH2  | CaH3  | CaH2+CaH3 |
|--------------|-------|-------|-------|-------|-----------|
| LHS0326      | 0.948 | 0.782 | 0.689 | 0.841 | 1.530     |
| LHS0327      | 0.995 | 1.004 | 1.025 | 1.005 | 2.030     |
| LHS0334      | 0.475 | 0.492 | 0.277 | 0.470 | 0.747     |
| LHS0335      | 0.960 | 0.721 | 0.640 | 0.791 | 1.431     |
| LHS0360      | 0.986 | 0.840 | 0.778 | 0.886 | 1.664     |
| LHS0367      | 1.011 | 0.880 | 0.854 | 0.929 | 1.783     |
| LHS0375      | 0.879 | 0.604 | 0.414 | 0.583 | 0.997     |
| LHS0381      | 0.883 | 0.728 | 0.555 | 0.730 | 1.285     |
| LHS0385      | 0.982 | 0.787 | 0.712 | 0.841 | 1.553     |
| LHS0398      | 0.910 | 0.767 | 0.644 | 0.808 | 1.452     |
| LHS0401      | 0.997 | 0.959 | 0.911 | 0.958 | 1.869     |
| LHS0406      | 0.686 | 0.801 | 0.576 | 0.789 | 1.365     |
| LHS0418      | 0.913 | 0.924 | 0.858 | 0.924 | 1.782     |
| LHS0424      | 0.951 | 0.917 | 0.852 | 0.927 | 1.778     |
| LHS0440      | 0.763 | 0.789 | 0.608 | 0.796 | 1.404     |
| LHS0507      | 0.991 | 0.958 | 0.969 | 0.980 | 1.949     |
| LHS0515      | 0.813 | 0.524 | 0.355 | 0.508 | 0.863     |
| LHS0518      | 0.966 | 0.933 | 0.903 | 0.951 | 1.854     |
| LHS0521      | 1.006 | 0.937 | 0.950 | 0.969 | 1.919     |
| LHS0541      | 0.795 | 0.678 | 0.538 | 0.701 | 1.239     |
| LHS1490      | 0.301 | 0.778 | 0.296 | 0.569 | 0.865     |
| LHS1970      | 0.878 | 0.433 | 0.476 | 0.686 | 1.162     |
| LHS2067A     | 0.228 | 0.670 | 0.199 | 0.452 | 0.651     |
| LHS2467      | 1.017 | 1.031 | 1.019 | 1.014 | 2.033     |
| LHS2734A     | 1.018 | 0.985 | 0.984 | 0.987 | 1.972     |
| LHS2734B     | 1.086 | 0.720 | 0.669 | 0.828 | 1.497     |
| LHS3620      | 0.836 | 0.684 | 0.544 | 0.727 | 1.271     |
| SCR0242–5935 | 0.910 | 0.906 | 0.838 | 0.891 | 1.728     |
| SCR0406–6735 | 1.015 | 0.989 | 0.829 | 0.937 | 1.766     |
| SCR0433–7740 | 0.929 | 0.887 | 0.760 | 0.864 | 1.624     |
| SCR0529–3950 | 0.817 | 0.886 | 0.637 | 0.815 | 1.452     |

Table 1—Continued

| Object       | TiO5  | CaH1  | CaH2  | CaH3  | CaH2+CaH3 |
|--------------|-------|-------|-------|-------|-----------|
| SCR0629–6938 | 0.845 | 0.908 | 0.579 | 0.784 | 1.363     |
| SCR0654–7358 | 0.938 | 0.835 | 0.653 | 0.825 | 1.478     |
| SCR0701–0655 | 0.976 | 0.886 | 0.780 | 0.881 | 1.662     |
| SCR0708–4709 | 0.984 | 0.992 | 0.979 | 0.976 | 1.955     |
| SCR0709–4648 | 1.013 | 0.899 | 0.846 | 0.915 | 1.761     |
| SCR1107–4135 | 0.994 | 0.839 | 0.783 | 0.892 | 1.675     |
| SCR1433–3847 | 1.001 | 0.904 | 0.797 | 0.930 | 1.726     |
| SCR1455–3914 | 0.915 | 0.788 | 0.707 | 0.849 | 1.555     |
| SCR1457–3904 | 0.905 | 0.824 | 0.707 | 0.838 | 1.545     |
| SCR1613–3040 | 0.920 | 0.898 | 0.749 | 0.885 | 1.634     |
| SCR1739–8222 | 0.945 | 1.009 | 0.797 | 0.882 | 1.679     |
| SCR1740–5646 | 0.915 | 0.696 | 0.496 | 0.637 | 1.133     |
| SCR1756–5927 | 0.922 | 0.794 | 0.757 | 0.856 | 1.613     |
| SCR1822–4542 | 0.817 | 0.818 | 0.624 | 0.809 | 1.433     |
| SCR1843–7849 | 0.869 | 0.810 | 0.715 | 0.829 | 1.545     |
| SCR1913–1001 | 0.962 | 0.923 | 0.835 | 0.925 | 1.760     |
| SCR1916–3638 | 0.931 | 0.647 | 0.493 | 0.666 | 1.159     |
| SCR1958–5609 | 0.915 | 0.865 | 0.805 | 0.926 | 1.731     |
| SCR2018–6606 | 0.886 | 0.856 | 0.715 | 0.836 | 1.550     |
| SCR2101–5437 | 0.929 | 0.969 | 0.726 | 0.904 | 1.630     |
| SCR2104–5229 | 0.940 | 0.907 | 0.772 | 0.865 | 1.638     |
| SCR2109–5226 | 0.967 | 0.774 | 0.646 | 0.799 | 1.445     |
| SCR2204–3347 | 0.874 | 0.711 | 0.569 | 0.762 | 1.331     |
| SIP1342–3534 | 0.684 | 0.664 | 0.462 | 0.676 | 1.138     |
| WT0135       | 0.622 | 0.754 | 0.512 | 0.731 | 1.243     |
| WT0233       | 0.939 | 0.933 | 0.739 | 0.857 | 1.596     |

Table 2. Spectral Types

| RA         | DEC         | Object       | $K_s$        | $V - K_s$ | $M_{K_s}$ | $V_{tan}$<br>km/sec | Old Type  | New Type | metallicity              | gravity | $\pi$ | Ref<br>V spect |
|------------|-------------|--------------|--------------|-----------|-----------|---------------------|-----------|----------|--------------------------|---------|-------|----------------|
| 12 06 00.9 | +14 38 56.8 | G012-016     | 7.931±0.018  | 2.16      | 4.12      | 98.4                | K2        | GVI      | (-0.52) <sup>a</sup>     |         | H     | 2 2            |
| 01 04 26.4 | -02 21 59.8 | G070-035     | 7.116±0.020  | 2.04      | 4.15      | 55.8                | G5        | GVI      | (-0.67) <sup>a</sup>     |         | H     | 4 H            |
| 02 25 49.8 | +05 53 39.5 | G073-056     | 10.427±0.023 | 2.06      | 8.64      | 44.9                |           | GVI      | (-1.18) <sup>b</sup>     |         | H     | 4              |
| 07 54 34.1 | -01 24 44.3 | G112-054     | 5.425±0.023  | 2.00      | 4.01      | 23.7                | K1        | GVI      | (-0.94) <sup>c</sup>     |         | H     | 1 2            |
| 16 13 48.6 | -57 34 13.8 | LHS0413      | 5.293±0.024  | 2.24      | 4.60      | 106.5               | G8/K0V(W) | GVI      | (-1.35) <sup>c</sup>     |         | H     | 1 10           |
| 15 45 52.4 | +05 02 26.6 | G016-009AB   | 6.880±0.024  | 2.27      | 3.46      | 59.2                | K2V       | K3.0[VI] | m-, (-0.77) <sup>d</sup> | g       | H     | H 3            |
| 00 50 17.0 | -39 30 08.3 | LHS0125      | 11.452±0.026 | 2.88      | 7.73      | 342.6               |           | K4.0[VI] | m-                       | g       | R     | R              |
| 07 35 46.3 | +03 29 36.0 | LHS0232      | 10.841±0.024 | 2.84      | 6.57      | 346.0               |           | K4.0[VI] | m-                       | g       | Y     | R              |
| 12 25 50.7 | -24 33 17.8 | LHS0327      | 10.144±0.021 | 2.57      | 5.44      | 418.3               | K0V       | K4.0[VI] | m-                       | g       | R     | R 2            |
| 11 52 32.0 | +27 30 51.3 | LHS2467      | 9.806±0.017  | 2.43      | 4.94      | 437.0               | G7V       | K4.0[VI] | m-                       | g       | H     | R 2            |
| 19 07 02.0 | +07 36 57.3 | G022-015     | 6.469±0.018  | 2.71      | 4.54      | 97.2                | K5V       | K5.0[VI] | m-, (-0.61) <sup>e</sup> | g       | H     | H 2            |
| 21 32 11.9 | +00 13 18.0 | G026-009ACD  | 7.082±0.029  | 2.64      | 3.62      | 96.9                | K2V       | K5.0[VI] | m-, (-1.05) <sup>f</sup> | g       | H     | H 2            |
| 05 54 34.1 | -09 23 33.7 | GJ0223.1     | 7.760±0.020  | 2.96      | 4.84      | 82.1                | K4V       | K5.0[VI] | m-, (-0.62) <sup>e</sup> | g       | H     | 1 9            |
| 04 32 36.6 | -39 02 03.4 | LHS0193A     | 8.427±0.023  | 3.23      | 5.80      | 147.3               |           | K6.0VI   | m-                       | g       | R     | 16 44          |
| 23 43 16.7 | -24 11 16.4 | LHS0073      | 9.393±0.021  | 3.42      | 7.27      | 322.3               | K5V       | K6.0VI   | m-                       | g+      | Y     | 19 2           |
| 07 13 40.6 | -13 27 57.1 | LHS0227      | 11.036±0.023 | 3.41      | 7.15      | 362.5               |           | K6.0VI   | m-                       | g+      | Y     | R              |
| 02 52 45.7 | +01 55 49.4 | LHS0161      | 10.995±0.019 | 3.65      | 8.01      | 272.6               | esdM2     | K6.0VI   | m-                       | g++     | Y     | R 8            |
| 03 01 40.6 | -34 57 56.5 | LHS0164      | 10.641±0.024 | 2.92      | 8.20      | 192.6               |           | K7.0VI   | m--                      | g       | R     | R              |
| 05 15 45.1 | -72 11 22.2 | DEN0515-7211 | 13.211±0.040 | 3.29      | 9.72      | 4.7                 |           | K7.0VI:  | m--                      | g       | 7     | 7              |
| 07 08 32.0 | -47 09 30.5 | SCR0708-4709 | 10.764±0.025 | 3.05      | ...       | ...                 |           | K7.0VI:  | m--                      | g       |       | R              |
| 13 25 14.0 | -21 27 06.0 | LHS2734A     | 12.896±0.037 | 3.23      | too far   | >280                |           | K7.0VI   | m-                       | g       | R     | R              |
| 03 06 28.7 | -07 40 41.5 | LHS0165      | 11.006±0.025 | 3.42      | 7.67      | 333.1               |           | M0.0VI   | m- - -                   | g+:     | Y     | R              |
| 11 11 13.7 | -41 05 32.7 | LHS0300A     | 9.802±0.023  | 3.38      | 7.23      | 197.5               |           | M0.0VI:  | m--                      | g       | 11    | 16             |
| 16 25 14.0 | +15 40 54.2 | LHS0418      | 10.072±0.018 | 3.37      | 6.44      | 301.5               | K7V       | M0.0VI   | m--                      | g       | Y     | R 8            |
| 16 37 05.6 | -01 32 01.6 | LHS0424      | 10.803±0.022 | 3.37      | 7.39      | 279.6               |           | M0.0VI   | m--                      | g       | Y     | R              |
| 08 13 27.8 | -09 27 56.6 | LHS0244      | 10.729±0.023 | 3.64      | 6.88      | 412.7               |           | M0.0VI   | m-                       | g++:    | Y     | 19             |
| 22 27 59.0 | -30 09 30.0 | LHS0521      | 11.463±0.019 | 3.22      | 8.14      | 222.1               |           | M0.5VI:  | m- - -                   | g       | 11    | 11             |
| 21 21 34.8 | -19 03 38.6 | LHS0507      | 12.053±0.026 | 3.15      | 7.81      | 355.2               | K/M sd    | M0.5VI   | m- - -                   | g+      | Y     | R Y            |
| 15 39 39.0 | -55 09 10.0 | LHS0401      | 9.407±0.019  | 3.31      | 7.33      | 142.2               |           | M0.5VI   | m- - -                   | g++     | Y     | R              |
| 02 42 26.3 | -59 35 01.6 | SCR0242-5935 | 12.783±0.031 | 3.40      | ...       | ...                 |           | M0.5VI   | m--                      | g       |       | R              |
| 07 09 37.2 | -46 48 58.8 | SCR0709-4648 | 11.491±0.026 | 3.41      | ...       | ...                 |           | M0.5VI   | m--                      | g       |       | R              |
| 11 07 55.8 | -41 35 52.7 | SCR1107-4135 | 11.474±0.019 | 3.49      | ...       | ...                 |           | M0.5VI   | m--                      | g       |       | R              |
| 17 39 45.4 | -82 22 02.2 | SCR1739-8222 | 12.190±0.026 | 3.48      | ...       | ...                 |           | M0.5VI   | m--                      | g       |       | R              |
| 02 02 52.2 | +05 42 21.0 | LHS0012      | 8.684±0.020  | 3.56      | 6.32      | 342.1               | sdM0      | M0.5VI   | m-                       | g       | Y     | 13 8           |

Table 2—Continued

| RA         | DEC         | Object       | $K_s$        | $V - K_s$ | $M_{K_s}$ | $V_{tan}$<br>km/sec | Old Type | New Type | metallicity                | gravity | $\pi$ | Ref<br>V spect |
|------------|-------------|--------------|--------------|-----------|-----------|---------------------|----------|----------|----------------------------|---------|-------|----------------|
| 11 11 22.6 | −06 31 56.4 | LHS0299      | 11.143±0.024 | 3.64      | 6.54      | 437.3               |          | M0.5VI:  | m−                         | g       | Y     | R              |
| 14 18 20.4 | −52 24 12.6 | LHS0367      | 9.786±0.019  | 3.41      | 6.26      | 268.6               |          | M0.5VI:  | m−                         | g       | 7     | 7              |
| 04 06 06.7 | −67 35 28.8 | SCR0406−6735 | 12.804±0.031 | 3.54      | ...       | ...                 |          | M0.5VI   | m−                         | g       |       | R              |
| 04 33 26.5 | −77 40 09.7 | SCR0433−7740 | 13.361±0.034 | 3.50      | ...       | ...                 |          | M0.5VI   | m−                         | g       |       | R              |
| 14 33 03.3 | −38 46 59.6 | SCR1433−3847 | 13.592±0.046 | 3.62      | ...       | ...                 |          | M0.5VI   | m−                         | g       |       | R              |
| 18 43 35.7 | −78 49 02.5 | SCR1843−7849 | 12.591±0.026 | 3.55      | ...       | ...                 |          | M0.5VI   | m−                         | g       |       | R              |
| 19 58 31.2 | −56 09 10.6 | SCR1958−5609 | 12.525±0.033 | 3.51      | ...       | ...                 |          | M0.5VI   | m−                         | g       |       | R              |
| 13 46 55.5 | +05 42 56.4 | LHS0360      | 11.662±0.023 | 3.50      | 6.73      | 524.0               |          | M0.5VI:  | m−                         | g+      | 7     | 7              |
| 22 20 27.0 | −24 21 49.3 | LHS0518      | 10.393±0.021 | 3.24      | 7.07      | 230.7               |          | M1.0VI   | m− − − − −                 | g       | 11    | 11             |
| 07 01 17.7 | −06 55 49.3 | SCR0701−0655 | 12.996±0.030 | ...       | ...       | ...                 |          | M1.0VI   | m− − − − −                 | g+      |       |                |
| 01 53 09.0 | −33 25 02.1 | LHS0148      | 12.832±0.032 | 3.59      | 8.59      | 374.8               |          | M1.0VI   | m− − − − −                 | g+:     | 6     | 6              |
| 01 53 09.0 | −33 25 02.1 | SCR1913−1001 | 11.929±0.028 | 3.69      | ...       | ...                 |          | M1.0VI   | m− − − −                   | g       |       | R              |
| 00 17 40.0 | −10 46 16.9 | LHS0109      | 10.366±0.021 | 3.51      | 7.63      | 176.1               | K5       | M1.0VI   | m− − − −                   | g+      | Y     | R              |
| 21 04 00.5 | −52 29 43.5 | SCR2104−5229 | 12.763±0.024 | 3.53      | ...       | ...                 |          | M1.0VI   | m− − − −                   | g+      |       | R              |
| 14 55 35.8 | −15 33 44.0 | LHS0385      | 11.062±0.023 | 3.55      | 7.61      | 403.4               | M0       | M1.0VI   | m− − − −                   | g++     | Y     | R              |
| 14 55 51.5 | −39 14 33.1 | SCR1455−3914 | 11.788±0.024 | 3.65      | ...       | ...                 |          | M1.0VI   | m− − − −                   | g++     |       | R              |
| 12 34 53.1 | +05 03 54.1 | LHS0335      | 13.000±0.030 | 3.60      | 9.74      | 247.2               |          | M1.0VI   | m− − − −                   | g+++    | Y     | R              |
| 21 09 02.5 | −52 26 17.8 | SCR2109−5226 | 13.049±0.036 | 3.66      | ...       | ...                 |          | M1.0VI   | m− − − −                   | g+++    |       | R              |
| 01 31 04.1 | −50 24 54.3 | LEHPM1628    | 13.459±0.038 | 3.70      | ...       | ...                 |          | M1.0VI   | m− − − −                   | g++++   |       | R              |
| 06 54 06.3 | −73 58 03.6 | SCR0654−7358 | 13.285±0.048 | ...       | ...       | ...                 |          | M1.0VI   | m− − −                     | g       |       |                |
| 14 57 49.0 | −39 04 51.4 | SCR1457−3904 | 12.984±0.030 | 3.69      | ...       | ...                 |          | M1.0VI   | m− − −                     | g       |       | R              |
| 16 13 53.5 | −30 40 58.4 | SCR1613−3040 | 12.383±0.031 | 3.66      | ...       | ...                 |          | M1.0VI   | m− − −                     | g       |       | R              |
| 17 56 27.9 | −59 27 18.2 | SCR1756−5927 | 12.686±0.030 | 3.61      | ...       | ...                 |          | M1.0VI   | m− − −                     | g       |       | R              |
| 20 18 28.7 | −66 06 44.5 | SCR2018−6606 | 12.990±0.030 | ...       | ...       | ...                 |          | M1.0VI   | m− − −                     | g       |       |                |
| 05 29 40.9 | −39 50 25.6 | SCR0529−3950 | 11.651±0.021 | ...       | ...       | ...                 |          | M1.0VI   | m− −                       | g       |       |                |
| 21 01 45.6 | −54 37 31.9 | SCR2101−5437 | 12.078±0.026 | 3.70      | ...       | ...                 |          | M1.0VI   | m−                         | g       |       | R              |
| 08 01 29.0 | +10 43 04.2 | LHS1970      | 13.875±0.032 | 3.84      | 9.43      | 374.8               | esdM2.5  | M1.0VI:  | m−                         | g+:     | Y     | R              |
| 13 25 14.0 | −21 27 06.0 | LHS2734B     | 14.932±0.136 | 3.90      | too far   | >280                |          | M1.0VI:  | m−                         | g+:     | R     | R              |
| 02 42 02.9 | −44 30 58.7 | LHS0158      | 9.726±0.021  | 3.91      | 6.89      | 191.4               |          | M1.0[VI] | m                          | g       | 11    | 11             |
| 17 18 35.0 | −43 26 24.0 | LHS0440      | 8.948±0.023  | 4.03      | 6.78      | 135.6               | M1V      | M1.0VI:  | m                          | g+:     | 11    | 11             |
| 18 22 58.7 | −45 42 45.3 | SCR1822−4542 | 12.879±0.027 | 3.92      | ...       | ...                 |          | M1.0VI:  | m                          | g+:     |       | R              |
| 07 56 13.4 | −67 05 20.6 | WT0233       | 12.627±0.024 | 3.60      | 7.84      | 326.8               | M0.0VI   | M2.0VI   | m− − −                     | g       | 6     | 6              |
| 11 56 54.8 | +26 39 56.3 | LHS0318      | 11.797±0.018 | 3.65      | 8.77      | 263.8               |          | M2.0VI:  | m− − −                     | g       | 18    | R              |
| 05 11 40.6 | −45 01 06.0 | GJ0191       | 5.049±0.021  | 3.80      | 7.08      | 161.0               | sdM1.0   | M2.0VI   | m− −, (−0.99) <sup>e</sup> | g       | Y     | 13             |

2  
−  
45  
−

Table 2—Continued

| RA         | DEC         | Object       | $K_s$        | $V - K_s$ | $M_{K_s}$ | $V_{tan}$<br>km/sec | Old Type | New Type | metallicity | gravity | $\pi$ | Ref |       |  |
|------------|-------------|--------------|--------------|-----------|-----------|---------------------|----------|----------|-------------|---------|-------|-----|-------|--|
|            |             |              |              |           |           |                     |          |          |             |         |       | V   | spect |  |
| 02 56 13.2 | −35 08 26.9 | LHS0162      | 11.536±0.021 | 3.82      | 7.13      | 364.1               | M1.0VI   | M2.0VI   | m−−         | g+      | 6     | 6   | 6     |  |
| 21 04 25.5 | −27 52 48.5 | LHS3620      | 12.696±0.027 | 3.93      | 7.94      | 414.6               |          | M2.0VI   | m−−         | g+      | R     | R   |       |  |
| 04 03 38.4 | −05 08 05.4 | LHS0186      | 10.854±0.024 | 4.02      | 7.21      | 295.6               |          | M2.0VI   | m−          | g       | Y     | R   |       |  |
| 15 43 18.3 | −20 15 33.0 | LHS0406      | 9.018±0.021  | 4.04      | 7.40      | 117.4               | M1.0V    | M2.0VI   | m−          | g       | 11    | 11  | 2     |  |
| 00 55 43.8 | −21 13 05.5 | LHS0127      | 11.733±0.023 | 4.06      | ...       | ...                 |          | M2.0VI   | m−          | g+      |       | R   |       |  |
| 06 29 56.4 | −69 38 13.4 | SCR0629−6938 | 12.901±0.034 | ...       | ...       | ...                 |          | M2.0VI   | m−          | g+      |       |     |       |  |
| 12 24 26.8 | −04 43 36.7 | LHS0326      | 11.234±0.023 | 3.67      | 7.69      | 304.9               |          | M3.0VI   | m−−         | g       | R     | R   |       |  |
| 15 34 27.7 | +02 16 47.5 | LHS0398      | 11.502±0.025 | 3.74      | 7.52      | 354.3               |          | M3.0VI   | m−−         | g       | Y     | 19  |       |  |
| 23 17 05.0 | −13 51 04.1 | LHS0541      | 12.414±0.026 | 4.05      | 8.22      | 420.0               | sdM3.0   | M3.0VI   | m−−         | g+      | Y     | 1   | 8     |  |
| 22 04 02.2 | −33 47 38.9 | SCR2204−3347 | 11.601±0.027 | 3.84      | ...       | ...                 |          | M3.0VI   | m−−         | g+      |       | R   |       |  |
| 17 40 46.9 | −56 46 58.0 | SCR1740−5646 | 13.195±0.040 | ...       | ...       | ...                 |          | M3.0VI   | m−−         | g++     |       |     |       |  |
| 19 16 46.5 | −36 38 05.8 | SCR1916−3638 | 12.947±0.034 | 3.88      | ...       | ...                 |          | M3.0VI   | m−−         | g++     |       | R   |       |  |
| 04 25 38.4 | −06 52 37.0 | LHS0189AB    | 10.311±0.037 | 3.94      | 8.59      | 128.3               |          | M3.0VI:  | m−          | g       | 7     | 7   |       |  |
| 07 16 27.7 | +23 42 10.4 | LHS0228      | 11.298±0.018 | 4.20      | 7.54      | 300.6               | sdM2.0   | M3.0VI   | m−          | g       | Y     | R   | 14    |  |
| 13 42 21.2 | −35 34 50.7 | SIP1342−3534 | 12.935±0.030 | 4.37      | ...       | ...                 |          | M3.0VI   | m−:         | g       |       | R   |       |  |
| 04 11 27.1 | −44 18 09.0 | WT0135       | 9.834±0.020  | 4.33      | ...       | ...                 |          | M3.0VI   | m−:         | g       |       | 5   |       |  |
| 09 43 46.2 | −17 47 06.2 | LHS0272      | 8.874±0.021  | 4.29      | 8.34      | 86.3                | sdM3.0   | M3.0VI   | m−:         | g+      | Y     | R   | 8     |  |
| 14 50 28.8 | −08 38 36.8 | LHS0381      | 11.237±0.021 | 3.90      | 8.43      | 274.9               | K7.0V    | M3.5VI   | m−−         | g       | Y     | 19  | Y     |  |
| 14 31 38.4 | −25 25 33.9 | LHS0375      | 11.507±0.022 | 4.12      | 9.61      | 157.5               | esdM4    | M3.5VI   | m−−         | g+      | Y     | R   | 8     |  |
| 01 38 49.0 | +11 21 36.7 | LHS0144      | 12.080±0.026 | 4.24      | 8.57      | 386.6               |          | M3.5VI   | m−          | g       | Y     | R   |       |  |
| 21 55 48.0 | −11 21 42.1 | LHS0515      | 12.912±0.032 | 4.50      | 9.36      | 264.5               | esdM5    | M3.5VI   | m−          | g+      | Y     | R   | 17    |  |
| 05 00 15.3 | −54 06 09.0 | LEHPM3861    | 13.967±0.059 | 4.47      | ...       | ...                 | esdM6    | M4.0VI   | m−          | g       |       | R   | 15    |  |
| 03 02 06.3 | −39 50 51.8 | LHS1490      | 9.885±0.023  | 4.37      | ...       | ...                 |          | M5.0VI   | m−          | g       |       | R   |       |  |
| 12 34 15.7 | +20 37 05.7 | LHS0334      | 13.044±0.029 | 4.98      | ...       | ...                 | sdM4.5   | M6.0VI   | m−−         | g       |       | R   | 17    |  |
| 08 53 57.0 | −24 46 54.0 | LHS2067A     | 11.571±0.023 | 6.38      | 9.54      | 76.2                | subdwarf | M6.0VI   | m−          | g       | R     | R   | 12    |  |

Note. — The last three columns provide references for parallaxes,  $V$  band photometry, and previous spectroscopy results. “:” indicates a questionable sub-type, metallicity, or gravity. All  $K_s$  magnitudes are from the 2MASS all-sky database. LHS2734A and B have zero parallax so we list them as “too far” in the table.

<sup>a</sup>Metallicity data are from Carney et al. (1994).

<sup>b</sup>Metallicity data are from Nordström et al. (2004).

<sup>c</sup>Metallicity data are from Cayrel de Strobel et al. (2001). There are multiple metallicities reported for G112-054, so a mean value is presented.

<sup>d</sup>Cayrel de Strobel et al. (2001), Goldberg et al. (2002) and Laird et al. (1988) report different metallicities, so a mean value is given.

<sup>e</sup>Metallicity data are from Woolf & Wallerstein (2005)

<sup>f</sup>Morrison et al. (2003) and Allen et al. (2000) report different metallicities, so a mean value is given.

References. — (1) Bessel 1990 ;(2) Bidelman 1985 ;(3) Cayrel de Strobel et al. 1997 ;(4) Carney et al. 1994 ;(5) Costa & Méndez 2003 ;(6) Costa et al. 2005 ;(7) Costa et al. 2006 ;(8) Gizis 1997 ;(9) Hawley et al. 1996 ;(10) Houk & Cowley 1975 ;(11) Jao et al. 2005 ;(12) Kirkpatrick et al. 1995 ;(13) Leggett 1992 ;(14)Lépine et al. 2007 ;(15) Lodieu et al. 2005 ;(16) Monteiro et al. 2006 ;(17) Reid and Gizis 2005 ;(18) Smart et al. 2007 ;(19) Weis 1996 ;(H) ESA 1997 ;(R) Jao in preparation 2008 ;(Y) van Altena et al. 1995

Table 3. SDSS Subdwarfs

| Object                   | $r$   | $g - r$ | $g - z$ | Old Type | New Type | metallicity | gravity |
|--------------------------|-------|---------|---------|----------|----------|-------------|---------|
| SDSS J085843.89+511210.1 | 20.00 | 1.53    | 3.10    | M2.0VI   | M2.0V    | m           | ...     |
| SDSS J093141.85+453914.5 | 19.45 | 1.58    | 3.08    | M2.0VI   | M2.0V    | m           | ...     |
| SDSS J145447.32+011006.8 | 20.35 | 1.44    | 3.45    | M3.0VI   | M3.0V    | m           | ...     |
| SDSS J003755.20-002134.2 | 18.70 |         |         | M1.0VI   | M2.0VI   | m-          | g       |
| SDSS J083217.77+522408.2 | 19.61 | 1.49    | 3.14    | M2.0VI   | M2.0VI:  | m           | g+:     |
| SDSS J033408.64-072349.2 | 20.21 | 1.89    | 3.35    | M2.0VI   | M3.0VI   | m- - -      | g+:     |
| SDSS J090434.02+513153.9 | 19.40 | 1.78    | 3.29    | M1.0VI   | M3.0VI   | m- - -      | g+:     |
| SDSS J161348.84+482016.0 | 18.30 | 1.72    | 3.19    | M1.0VI   | M3.0VI   | m- - -      | g+:     |
| SDSS J112751.35-001246.8 | 20.03 |         |         | M1.0VI   | M3.0VI   | m- - -      | g       |
| SDSS J092708.10+561648.1 | 19.37 | 1.62    | 3.02    | M1.0VI   | M3.0VI   | m- - -      | g+      |
| SDSS J100109.54+015450.2 | 19.12 | 1.71    | 3.25    | M2.0VI   | M3.0VI   | m- - -      | g++     |
| SDSS J101031.13+651327.6 | 19.39 | 1.68    | 3.30    | M1.0VI   | M3.0VI   | m- - -      | g++     |
| SDSS J002228.00-091444.8 | 18.92 | -0.89   | 3.03    | M1.0VI   | M3.0VI   | m--         | g       |
| SDSS J024501.77+003315.8 | 19.38 | 1.54    | 3.15    | M1.0VI   | M3.0VI   | m--         | g       |
| SDSS J081329.95+443945.6 | 19.39 | 1.68    | 3.35    | M2.0VI   | M3.0VI   | m--         | g       |
| SDSS J090238.75+471813.6 | 19.97 | 1.73    | 3.21    | M2.0VI   | M3.0VI   | m--         | g       |
| SDSS J092534.16+524442.4 | 19.79 | 1.71    | 3.31    | M3.0VI   | M3.0VI   | m--         | g       |
| SDSS J092745.78+582122.7 | 20.47 | 1.78    | 3.56    | M3.0VI   | M3.0VI   | m--         | g       |
| SDSS J095147.77+003612.0 | 18.27 | 1.58    | 3.15    | M2.0VI   | M3.0VI   | m--         | g       |
| SDSS J115900.70+665214.3 | 19.37 | 1.61    | 3.35    | M3.0VI   | M3.0VI   | m--         | g       |
| SDSS J125919.29-025402.3 |       |         |         | M2.0VI   | M3.0VI   | m--         | g       |
| SDSS J173452.52+603603.1 | 18.78 | 1.71    | 3.36    | M2.0VI   | M3.0VI   | m--         | g       |
| SDSS J215937.69+005536.2 | 18.96 | 1.61    | 3.22    | M2.0VI   | M3.0VI   | m--         | g       |
| SDSS J221500.88+005217.2 | 19.08 | 1.60    | 3.33    | M2.0VI   | M3.0VI   | m--         | g       |
| SDSS J145547.00+602837.3 | 19.18 | 1.70    | 3.39    | M2.0VI   | M3.0VI   | m--         | g+      |
| SDSS J104320.47+010439.4 | 19.16 | 1.66    | 3.37    | M2.0VI   | M3.0VI   | m--         | g++     |
| SDSS J084105.39+032109.6 | 20.05 | 1.45    | 3.18    | M3.0VI   | M3.0VI   | m-          | g       |
| SDSS J091451.98+453152.8 | 19.02 | 1.64    | 3.49    | M3.0VI   | M3.0VI   | m-          | g       |
| SDSS J093024.66+554447.7 | 19.14 | 1.56    | 2.96    | M3.0VI   | M3.0VI   | m-          | g       |
| SDSS J094306.37+465701.4 | 19.74 | 1.59    | 3.29    | M3.0VI   | M3.0VI   | m-          | g       |
| SDSS J224854.83-091723.2 | 19.85 | 1.57    | 3.39    | M3.0VI   | M3.0VI   | m-          | g       |



Table 3—Continued

| Object                   | $r$   | $g - r$ | $g - z$ | Old Type | New Type | metallicity | gravity |
|--------------------------|-------|---------|---------|----------|----------|-------------|---------|
| SDSS J235830.60–011413.2 | 19.99 | 1.54    | 3.33    | M2.0VI   | M3.0VI   | m–          | g       |
| SDSS J113501.76+033720.3 |       |         |         | M3.0VI   | M3.0VI:  | m           | g+:     |
| SDSS J105122.43+603844.8 | 17.15 | 1.65    | 3.21    | M2.0VI   | M4.0VI   | m– – –      | g       |
| SDSS J092429.76+523410.7 | 18.70 | 1.67    | 3.21    | M2.0VI   | M4.0VI   | m– – –      | g       |
| SDSS J031314.28–000619.8 | 20.33 | 1.48    | 3.39    | M2.0VI   | M4.0VI   | m– –        | g       |
| SDSS J082230.00+471645.8 | 19.39 | 1.67    | 3.38    | M2.0VI   | M4.0VI   | m– –        | g       |
| SDSS J143930.77+033317.3 | 19.38 | 1.80    | 3.58    | M3.0VI   | M4.0VI   | m– –        | g+      |
| SDSS J003541.84+003210.1 | 20.15 | 1.64    | 3.37    | M3.0VI   | M4.0VI   | m–          | g       |
| SDSS J003701.37–003248.3 | 20.15 | 1.60    | 3.50    | M3.0VI   | M4.0VI   | m–          | g       |
| SDSS J010811.89+003042.4 | 17.34 | 1.61    | 3.43    | M3.0VI   | M4.0VI   | m–          | g       |
| SDSS J083002.73+483251.6 | 19.92 | 1.90    | 3.69    | M3.0VI   | M4.0VI   | m–          | g       |
| SDSS J171745.22+625337.0 | 18.66 | 1.65    | 3.57    | M3.0VI   | M4.0VI   | m–          | g       |
| SDSS J221625.03–003122.5 | 19.28 | 1.58    | 3.37    | M3.0VI   | M4.0VI   | m–          | g       |
| SDSS J223802.82–082532.4 | 19.71 | 1.78    | 3.59    | M3.0VI   | M4.0VI   | m–          | g       |
| SDSS J224605.41+141640.6 | 17.00 | 1.57    | 3.48    | M3.0VI   | M4.0VI   | m–          | g       |
| SDSS J230303.49–010656.7 | 18.95 | 1.62    | 3.42    | M3.0VI   | M4.0VI   | m–          | g       |
| SDSS J235116.25–003104.8 | 19.48 | 1.49    | 3.31    | M3.0VI   | M4.0VI   | m–          | g       |
| SDSS J150511.33+620926.3 | 18.61 | 1.81    | 3.45    | M3.0VI   | M5.0VI   | m– –        | g       |
| SDSS J230805.24+001812.7 | 19.36 | 1.70    | 3.72    | M3.0VI   | M5.0VI   | m–          | g       |

Note. — Column definitions are the same as in Table 2. Double lines separate each type.




Fig. 1.— Synthetic spectra for stars having  $T_{eff} = 3500\text{K}$  and  $\log g = 5.0$ . Red, green, and blue lines represent different metallicities, 0.0,  $-2.0$  and  $-4.0$ . Note that the relative amounts of blue and red fluxes trends toward bluer objects at lower metallicities.




Fig. 2.— The mass-gravity relation is shown using data from Table 1 of López-Morales (2007). A solid line represents a polynomial fit to all of the points, simply to be used as a guide. Different shades for the points represent different metallicity measurements. Stars with unknown metallicities are plotted as open circles.




Fig. 3.— The CaH1 and CaH2+CaH3 versus TiO5 indices are plotted for our identified subdwarfs (solid circles). For comparison, known cool dwarfs (dots), subdwarfs (open triangles), and “extreme” subdwarfs (open boxes) from Hawley et al. (1996), Gizis (1997), and Reid and Gizis (2005) are also shown. The dashed box indicates subdwarfs that do not exhibit strong spectroscopic indices, but which are either below the main-sequence on the HR diagram or have published metallicities  $[m/H]$  less than or equal to  $-0.5$ . A solid line indicates the separation between regular subdwarfs and extreme subdwarfs adopted by Gizis (1997).




Fig. 4.— HR diagram for subdwarfs listed in Table 1 and shown in Figure 3 that have trigonometric parallaxes. Open circles indicate subdwarfs and filled boxes indicate “extreme” subdwarfs based on spectroscopic indices. Concentric circles indicate stars with CaH1 index greater than 0.9 that are difficult to distinguish from main sequence stars at our spectral resolution. Open boxes indicate confirmed spectroscopic binaries. A solid line indicates a fit to main-sequence dwarfs, primarily from Henry et al. (2004) with extra dwarf standard stars from Gray et al. (2003). The dashed line is one magnitude fainter than this solid line. Note that the K-type subdwarf sequence merges with the K dwarf sequence at the blue end of this ( $M_{K_s}$  vs  $V - K_s$ ) plot. The single point at  $V - K_s = 6.4$  is LHS 2067A. The spectral types for dwarfs are given at the bottom of the figure as references.




Fig. 5.— Spectra for G and K-type stars from Jacoby et al. (1984) (left panel,  $4\text{\AA}$  resolution) and Silva & Cornell (1992) (right panel,  $11\text{\AA}$  resolution) are shown. Because the red cutoff is  $7400\text{\AA}$  in Jacoby et al. (1984), the results from Silva & Cornell (1992) are also plotted to  $7400\text{\AA}$ , and both sets of spectra are normalized at  $7400\text{\AA}$ . The two tick marks indicate the Ba I (left) and  $H\alpha$  (right) absorption features. Silva & Cornell (1992) have removed telluric absorption features.




Fig. 6.— Top: An mid K subdwarf spectrum (black) is virtually identical to an mid K dwarf (red). Middle: The spectra of two early M subdwarfs differ only at CaH. Bottom: One subdwarf’s spectrum (black) matches an M0.5V spectral standard (red) at the blue end, but matches an M2.0V spectral standard (red) at the red end. The deep telluric band, O<sub>2</sub>A (7570Å–7700Å) has been removed.




Fig. 7.— Two synthetic spectra from old (red) and new (black) model grids are shown. The physical parameters for both models are  $T_{eff} = 3600\text{K}$ ,  $\log g = 5.0$  and  $[m/H] = -2.0$ . Major differences are seen between  $6500\text{\AA}$  and  $7000\text{\AA}$ , and in the absorption lines of Li I ( $6103\text{\AA}$ ) and Ca I ( $6122\text{\AA}$  and  $6162\text{\AA}$ ).






Fig. 8.— CaH1 (top) and CaH2+CaH3 (bottom) indices from GAIA model grids are plotted against  $T_{\text{eff}}$ . The different line styles represent different  $[m/H]$ . The spectral types for dwarfs are given at the top of the figure as references.




Fig. 9.— GAIA synthetic spectra from 4800K to 2800K are shown. All spectra are noiseless and have  $\log g = 5.0$ . Red, green and blue lines represent  $[m/H] = 0.0, -1.0$  and  $-2.0$ , respectively. Effective temperatures for each set of spectra are given above each group of lines. The feature marked #1 is not seen in any of our spectra. The Ca I (6162Å) feature marked #2 is seen in our spectra.




Fig. 10.— The top plot shows GAIA model grids at fixed metallicity ( $[m/H] = -1.0$ ) and effective temperature (3500K). Black, red, green and yellow lines represent various  $\log g = 4.0, 4.5, 5.0$  and  $5.5$ , respectively. It is clear that CaH bands will be affected by changing gravity but TiO5 is not. The bottom plot shows model grids at fixed  $\log g = 5.0$  and  $T = 3500$ K. Black, red and green lines represent various  $[m/H] = 0.0, -1.0$  and  $-2.0$ , respectively. Note that model grids do not have telluric lines.




Fig. 11.— Using our spectra and metallicities measured independently by others, three different types of M dwarfs show the metallicity trend predicted by GAIA models. The gray lines represent lower metallicity stars in each pair. Metallicities from Bonfils et al. (2005) are given in each panel for the stars.




Fig. 12.— K3.0[VI] to K5.0[VI] spectra are shown compared to K dwarf standard spectra (red). The two telluric bands ( $O_2$  A and B) have been removed for clarity. The three thick tick-marks represent the locations of Ca I ( $6162\text{\AA}$ ) and Ca II ( $8542\text{\AA}$  and  $8662\text{\AA}$ ) absorption features. The inset plot shows the locations of the stars on the HR diagram, where filled circles represent subdwarfs and triangles represent main sequence standards of types K3.0V, K4.0V, and K5.0V (left to right). For clarity, the axis labels for the inset plot are not shown but are always  $V - K_s$  vs.  $M_{K_s}$ . The errors in absolute magnitudes are shown. However, because the errors for  $V - K_s$  are equal to or smaller than the filled circles, they are not shown. The solid line represents a fitted main sequence line and the dashed line is one magnitude fainter than the solid line.




Fig. 13.— Spectra of LHS 125, LHS 232 and LHS 327 (all K4.0[VI]), are shown. Note that the blue end of these spectra are at 6500Å because of problems with these particular spectra between 6000Å and 6500Å. The two telluric O<sub>2</sub> A and B bands have been removed. The triangle in the inset plot represents a K4.0V standard star. Symbols and lines have the same meanings as in Figure 12.




Fig. 14.— Spectra of LHS 193A, LHS 227 and LHS 161 (all K6.0VI), are shown with our K5.0V and K7.0V standard spectra. The telluric O<sub>2</sub> A band has been removed. The thick arrows in the HR diagram point toward sequentially higher gravity subdwarfs. Symbols and lines have the same meanings as in Figure 12.




Fig. 15.— Spectra of LHS 164, LHS 2734A (both K7.0VI), DEN 0515–7211, and SCR0708–4709 (both K7.0VI:) are shown, with our K7.0V standard spectrum (represented by a triangle in the inset plot). The telluric O<sub>2</sub> A band has been removed. Symbols and lines have the same meanings as in Figure 12.






Fig. 16.— Spectra of LHS 165, LHS 418, LHS 244 (all M0.0VI), and LHS 300AB (M0.0VI:) are shown with our M0.0V standard spectrum (represented by a triangle in the inset plot). The telluric O<sub>2</sub> A band has been removed. The hollow arrow indicates the shift on the HR diagram caused by decreasing metallicity. Symbols and lines have the same meanings as in Figure 12.




Fig. 17.— Spectra of LHS 507, SCR 0709-4648, LHS 12, LHS 401 (all M0.5VI), and LHS 521 (M0.5VI:) are shown with our M0.5V standard spectrum (represented by a triangle in the inset plot). The telluric O<sub>2</sub> A band has been removed. The hollow arrow indicates the shift on the HR diagram caused by decreasing metallicity. The solid arrow indicates the shift caused by higher gravity. Symbols and lines have the same meanings as in Figure 12.




Fig. 18.— Spectra of LHS 367, LHS 299, and LHS 360, are shown (all M0.5VI:). The telluric O<sub>2</sub> A band has been removed.




Fig. 19.— Progressions in metallicity and gravity effects in M1.0VI subdwarf spectra are shown. The top set of spectra illustrates spectra with metallicities increasing from black to red (top to bottom, metallicity scale from m- - - - - to m). The red line is our M0.5V standard spectrum (represented by a triangle in the inset plot). The bottom two sets of spectra illustrate gravity effects for metallicity scales m- - - - - and m- - - - -. The telluric O<sub>2</sub> A band has been removed. Symbols and lines have the same meanings as in Figure 17.




Fig. 20.— The top set of spectra illustrates gravity effects for metallicity scale  $m - - -$ . Spectra of LHS 440, LHS 1970, LHS LHS 2734B (all M1.0VI), and LHS 158 (M1.0[VI]), are also shown compared individually to our M1.0V standard. The telluric O<sub>2</sub> A band has been removed.




Fig. 21.— Spectra for WT 233, GJ 191, LHS 406, LHS 3620, LHS 127 (all M2.0VI) and LHS 318 (M2.0VI:) are shown with our M2.0V standard spectrum (represented by a triangle in the inset plot). The telluric O<sub>2</sub> A band has been removed. Symbols and lines have the same meanings as in Figure 17.




Fig. 22.— Spectra for six M3.0VI are shown, with our M3.0V standard spectrum in the top set of spectra (represented by a triangle in the inset plot). LHS 326 and LHS 228 show metallicity changes, while LHS 326, SCR2204–3347 and SCR1916–3638 show effects of gravity. WT 135 and LHS 272 are M3.0VI:. Note that the K I lines at 7665Å and 7699Å have appeared and blended in the O<sub>2</sub> A band. Symbols and lines have the same meanings as in Figure 17.




Fig. 23.— Spectra for LHS 381 and LHS 144 (both M3.5VI) are shown with our M3.5V standard spectrum (represented by a triangle in the inset plot). These three stars show a dramatic metallicity sequence for M3.5. The spectrum of LHS 228 (M3.0VI) is compared to that of LHS 381 to illustrate the slope difference between M3.0VI and M3.5VI (8200Å–9000Å). LHS 375 and LHS 515 have higher gravities than LHS 381 and LHS 144, respectively. Note that K I lines (7665Å and 7699Å) have appeared and blended in the O<sub>2</sub> A band. Symbols and lines have the same meanings as in Figure 17.






Fig. 24.— LEHPM 3861 (M4.0VI), LHS 1490 (M5.0VI), LHS 334 (M6.0VI), and LHS2067A (M6.0VI) subdwarf spectra are shown compared to our standard M3.0-7.0V standards. Only LHS2067A and the M6.0V standard (filled triangle) are shown in the HR diagram. Vertical dotted lines represent the three peaks we use to assist us to assign spectral types. The K I lines at 7665Å and 7699Å are clearly seen blended in the O<sub>2</sub> A band. Symbols and lines have the same meanings as in Figure 17.




Fig. 25.— Spectra are shown for subdwarfs from West et al. (2004) separated into types assigned by us by matching standard spectra (red) from Bochanski et al. (2007) in the region 8200Å–9000Å. Other colors (black, blue, green, and yellow) represent different metallicities at each type. Metallicity increases from black to red.




Fig. 26.— These plots indicate the gravity effects for SDSS subdwarfs of type M3VI and M4VI. The black, blue and green lines represent increasing gravities: g, g+ and g++.




Fig. 27.— Observed spectra of dwarf standard stars (red lines) compared with the best fitting synthetic spectra (black lines). Telluric lines are not present in the synthetic spectra. Six regions labeled from 1 to 6 are discussed in the text.




Fig. 28.— Synthetic models were compared to the observed spectrum of our M1.0V standard. The resulting  $\chi^2$  values are plotted against temperature for models with five different metallicities, at four different gravities each. The  $[m/H] = -0.5$  plot has the tightest curves at different  $\log g$ .




Fig. 29.— Comparisons of observed spectra for M1.0V (top) and M3.0V (bottom) are shown against the two best fitting synthetic spectra for each. The red line is the best fit, where  $T_{eff}/[m/H]/\log g = 3600/-0.5/4.5$  for M1.0, and  $3200/-0.5/4.5$  for M3.0. The blue line is the second best fit, where the values are  $3400/-1.0/4.0$  for M1.0 and  $3300/0.5/5.5$  for M3.0. Telluric bands are marked as gray boxes.




Fig. 30.— The best  $\chi^2$  values are shown for synthetic spectra fits to observed spectra for stars of types M0.0V to M5.5V (labeled relative to each type). The dashed line indicates the selected limit for reliability of fitting model spectra to observed spectra — above this line fits are deemed unreliable.




Fig. 31.— The spectrum of the subdwarf LHS 335 (black) is shown with the corresponding best fitting synthetic spectrum (red).






Fig. 32.— The CaH2+CaH3 versus TiO5 indices are plotted for subdwarfs discussed in this paper (filled circles). For comparison, known cool dwarfs (dots), subdwarfs (open triangles), and “extreme” subdwarfs (open boxes) from Hawley et al. (1996), Gizis (1997), and Reid and Gizis (2005) are also shown. Green circles are subdwarfs with the highest gravities at a given type (g with most +) in Table 2. The two blue circles represent LHS 400 and SCR 1822–4542, which have higher gravities than M dwarfs. Red circles represent the lowest metallicity stars in Table 2 at a given type (m with most –). Solids lines indicate pairs of stars having the same number of – in metallicity, but different gravities. A dotted line connects LHS 2734A and LHS 2734B, which is the only wide CPM binary in our sample. For comparison, the inset plot illustrates the indices (TiO5 vs. CaH2+CaH3) calculated from GAIA models for temperatures 2800–4400K. Axis labels are omitted, but both axes have the same ranges as the main figure. Three different metallicities, 0.0, –1.0 and –2.0, and two gravities,  $\log g=4.5$  (solid line) and 5.5 (dash line), are shown. Connections are drawn for stars with the same temperatures but different gravities.




Fig. 33.— CaH1 and TiO5 indices from GAIA model grids are plotted against effective temperature. The black, blue, red and green lines indicate  $\log g=4.0, 4.5, 5.0, 5.5$ , respectively. Note that a given CaH1 and TiO5 value may correspond to many combinations of metallicities and gravities.




Fig. 34.— Spectra from Lépine et al. (2007) are shown, with their published types. The colors (black and blue) represent different spectra, and their colors match with labels. Each pair has the almost the same overall spectral shape, but each star is assigned a different sub-type and identification as extreme or ultra. Spectra are normalized at 7500Å.




Fig. 35.— Spectra from Lépine et al. (2007) are shown for spectral types for M3.0, M5.0 and M7.5. The black, blue and red spectra represent sdM, esdM and usdM, respectively. Spectra are normalized at 7500Å.